

Index

Page references followed by f denote figures; those followed by t denote tables.

A

- A23187 (Ca²⁺ ionophore), 215, 216
 Acetamidiminodiacetic acid (ADA), affinity for Ca²⁺, 537t
 Acetoxymethyl (AM) esters of Ca²⁺ indicators, 408.
See also specific indicators
- Acids
 disposal of, 573
 safe handling of, 572, 575
 Acridine orange, 521–523
 ACSF for calcium imaging (recipe), 149, 152
 Action potentials, back propagating, 142, 142f
 Activation of Ca²⁺-dependent ion channels, 79–84
 ADA (acetamidiminodiacetic acid),
 affinity for Ca²⁺, 537t
 Adenophostin A, 173–174, 173f, 174f
 Adenosine diphosphoribose (ADPR), 498
 ADP, affinity for Ca²⁺, 537t
 ADP-ribosyl cyclase, 490, 493, 498–501, 518, 530
 Aequorin, 177–210, 193, 465–466
 advantages of, 180, 183, 194
 AquaLite, 203
 calibration curve, 179, 187, 188f
 as Ca²⁺-sensitive photoprotein, 178
 chimeras, 181t–182t
 decline in use as cytosolic probe, 179
 disadvantages of, 183
 experimental applications, 179–183
 f-aequorin, 203
 hcp-aequorin, 203
 history of, 177–178
 introduction into zebrafish embryos,
 193–210, 195f
 monitoring yeast intracellular Ca²⁺ levels using
 in vivo bioluminescence assay
 (protocol), 456–459
 materials, 456–457
 method, 457–459, 458f
 troubleshooting, 459
- protocols
 microinjecting holo-aequorin into
 dechorionated and intact zebrafish
 embryos, 197–205
 dechorionating embryos, 200
 discussion, 203
 imaging chamber preparation,
 198–200, 200f
 materials, 197–198
 method, 198–202
 micropipette preparation, 198, 199f
 recipes, 205
 troubleshooting, 202–203
 reconstitution of holo-aequorin with
 apoaequorin mRNA and
 coelenterazine in zebrafish
 embryos, 206–210
 discussion, 209
 materials, 206–207
 method, 207–209
 recipes, 210
 troubleshooting, 209
 using targeted variants of aequorin to
 measure Ca²⁺ levels in intracellular
 organelles, 185–191
 calibration, 187, 188f
 discussion, 191
 materials, 185–187, 186f
 measurement of Ca²⁺ in ER and its
 release, 190
 measurement of capacitative
 Ca²⁺ influx, 188–189
 measurement of cytosolic Ca²⁺,
 187–188
 measurement of mitochondrial
 Ca²⁺ uptake, 189, 189f
 measurement of mitochondrial Ca²⁺
 uptake in permeabilized cells, 190
 recipes, 191
 reconstitution of functional
 photoprotein, 187
 transfection, 187
 troubleshooting, 190–191
 reduction of affinity to measure high [Ca²⁺], 180
 semisynthetic, 193
 structure and function, 178–179, 178f, 193
 targeting of, 179–180, 181t–182t
 Aequorin buffer (recipe), 204
- Affinity
 of indicators for Ca²⁺, 28, 533–534
 of synthetic Ca²⁺ buffers, 537t
- Agarose, reactive red (RRA), 345
 Agarose solution for building agar salt bridges
 (recipe), 353
- Agatoxin IVA, 144
 Airy rings, 6
 Alcohol dehydrogenase, 495
 Alkaline phosphatase, 527
 Allen and Blinks algorithm, 187
 Alzheimer's disease, 323
 American Type Culture Collection (ATCC), 445, 575
 2-aminoethoxydiphenylborat (2-APB), 279, 287
 Amiodarone, 451
 AMP-5, affinity for Ca²⁺, 537t
 Amphotericin-B, 280
 Animals, humane treatment of, 573
 Antimycin, 215
 AP5, 144
 2-APB (2-aminoethoxydiphenylborat), 279, 287
 Apoaequorin, 193, 194, 206–209
 Apoptosis, 451–452
 AquaLite aequorin, 203
- Arabidopsis*
 high-resolution imaging of cytoplasmic Ca²⁺
 dynamics in *Arabidopsis* roots
 (protocol), 474–478
 discussion, 477
 materials, 474–475
 method, 475–477, 476f
 data analysis, 477
 imaging, 475–477
 plant growth, 475
 sample preparation, 475, 476f
 troubleshooting, 477
 imaging of mitochondrial and nuclear Ca²⁺
 dynamics in *Arabidopsis* roots
 (protocol), 484–488
 discussion, 487
 materials, 484–485
 method, 485–486
 CLSM analysis, 486
 imaging analyses, 486
 sample preparation, 486
 seed sterilization and growth of
 plantlets, 485–486
 recipes, 487
 troubleshooting, 487
- live cell imaging of cytoplasmic and nuclear
 Ca²⁺ dynamics in *Arabidopsis* roots
 (protocol), 479–483
 discussion, 482
 materials, 479–480
 method, 480–482, 481f
 Ca²⁺ imaging, 480
 data analyses, 482
 plant growth, 480
 sample preparation, 480, 481f
 recipes, 483
 troubleshooting, 482
- live cell imaging of cytoplasmic Ca²⁺ dynamics
 in *Arabidopsis* guard cells
 (protocol), 469–473
 discussion, 473
 materials, 469–470
 method, 470–472, 471f
 data analysis, 472
 imaging, 471–472
 plant growth, 470
 sample preparation, 470–471, 471f
 recipe, 473
 troubleshooting, 472–473
- Artificial seawater (ASW) (recipe), 515, 523
 Aspiration system, 38
 Astrocytes, intercellular Ca²⁺ waves in, 95, 96
 ATCC (American Type Culture Collection),
 445, 575
- ATP, 188
 affinity for Ca²⁺, 537t
 caged, 75, 111
 intercellular Ca²⁺ waves (ICWs) and, 95
- ATPases. *See also* Calcium (Ca²⁺) pump
 calculation of activity, 238, 239f
 plasma membrane Ca²⁺ ATPase (PMCA), 231,
 277, 291, 293, 438, 446
 secretory pathway Ca²⁺ ATPase (SPCA), 231, 446
 SERCA (*see* Sarco-endoplasmic Ca²⁺ ATPases)
 in yeast, 446
- ATP-regenerating system, 513, 522
 ATP stock solution for ER sensor calibrations
 (recipe), 63
- Autoclave safety, 572
 Auto-fluorescence, in two-photon microscopy, 16

B

- Background image, 34, 49
 Background intensity value, 34
 Background signal, subtraction of, 34, 35, 35f, 36f, 39
 Back-propagating action potentials (bAPs), 142, 142f
 Bacteria
 biological safety procedures, 574–575
 shipping requirements, 575

- BAPs (back-propagating action potentials), 142, 142f
- BAPTA
affinity for Ca^{2+} , 537t
 Ca^{2+} store depletion by, 279, 286
- BAPTA-dextran, 100
- Bases
disposal of, 573
safe handling of, 572, 575
- Bath solution for InsP_3R (recipe), 317
- Bath solution for recording Cl^- currents (recipe), 83, 88
- Bath solution for recording K^+ currents (recipe), 83, 88
- Bath solution for RyR (recipe), 318
- Bax inhibitor-1, 222
- BBB (blood-brain barrier), 95–96
- BCA test, 249
- Bcl-2, 222
- BDM (2,3-butanedione monoxime), 434
- Beclin 1, 222
- Berovin, 177
- Bicuculline, 144
- Binding rate constants of Ca^{2+} indicators, measuring, 541–545, 542t, 543f
- Biological safety procedures, 574–575
- Bioluminescence assay, monitoring yeast intracellular Ca^{2+} levels using in vivo (protocol), 456–459
materials, 456–457
method, 457–459, 458f
troubleshooting, 459
- Bis-Fura-2, dissociation constant of, 141, 142
- Bit depth, 31–32
- Black lipid membrane (BLM)
experiments to study InsP_3R and RyanR, 322–323
preparation of microsomes to study Ca^{2+} channels (protocol), 326–329
reconstitution of endoplasmic reticulum InsP_3 receptors into BLMs (protocol), 330–336
materials, 330–331, 331f
method, 331–334, 332f, 334f, 335f
recipes, 336
troubleshooting, 334–335
- Blebbistatin, 131
- Blocking filters, 4–5
- Blood-brain barrier (BBB), 95–96
- Blue light, 2
- Bradykinin, 96, 188
- Bradykinin-triggered Ca^{2+} oscillations, 99, 100
- BTP2, 287
- Buffer(s). *See also specific buffers*
affinity of synthetic Ca^{2+} buffers, 537t
effect on Ca^{2+} dynamics, 557
 pK_a and temperature dependent of common, 542t
- Buffer A for assaying Ca^{2+} transport by VDAC1 reconstituted into liposomes (recipe), 357
- Buffer A for isolation of mitochondria from rat liver (recipe), 346
- Buffer B for isolation of mitochondria from rat liver (recipe), 346
- Buffer B for microinjection (recipe), 507
- Buffer C for purification of VDAC1 from mitochondria (recipe), 347
- Buffer D for purification of VDAC1 from mitochondria (recipe), 347
- Buffer E for purification of VDAC1 from mitochondria (recipe), 347
- Buffer F for purification of VDAC1 from mitochondria (recipe), 347
- Buffer for assaying Ca^{2+} transport by VDAC1 reconstituted into PLBs (recipe), 353
- Buffer G for purification of VDAC1 from mitochondria (recipe), 347
- Buffer H for purification of VDAC1 from mitochondria (recipe), 347
- Buffer I for purification of VDAC1 from mitochondria (recipe), 348
- 2,3-butanedione monoxime (BDM), 434
- C**
- Cabuf program, 215
- Ca^{2+} -containing modified Krebs solution (recipe), 228
- cADPR. *See* Cyclic adenosine dinucleotide phosphate ribose
- CaEGTA (100 mM) (recipe), 436
- Ca-EGTA solution (recipe), 336
- Caffeine, 212, 425–428, 427f, 433, 433f, 442
- Ca^{2+} -free cytosol-like medium (recipe), 165
- Ca^{2+} -free perfusion solution (recipe), 124
- Caged Ca^{2+}
in Ca^{2+} -binding kinetics studies, 534–535
collecting data to determine the Ca^{2+} -binding properties of DM-nitrophen and proteins (protocol), 546–551, 547f
measuring rate constants of Ca^{2+} indicators (protocol), 541–545, 542t, 543f
in two-photon photolysis combined with kilobeam array scanner to probe calcium signaling in cardiomyocytes (protocol), 116–120
- Caged compounds, 75–92
applications, 76
caged NAADP (protocol), 525–528
in combining Ca^{2+} imaging with L-glutamate photorelease (protocol), 143–144, 144f, 150–152, 151f
cyclic adenosine dinucleotide phosphate ribose (cADPR), 528
inositol 1,4,5-trisphosphate (IP_3), 79, 81f, 90–92, 96, 99–101, 100f, 109–112, 528
intercellular Ca^{2+} waves (ICWs) and, 96
intracellular introduction of, 76–77
in nuclear microinjection to assess impact of heterologously expressed proteins on Ca^{2+} signals in *Xenopus* oocytes (protocol), 388–394
overview, 75–76
photolysis of
analyzing Ca^{2+} dynamics in intact epithelial cells using spatially limited flash photolysis (protocol), 90–92, 91f
materials, 90–91
method, 91–92, 91f
applications, 76
investigating ion channel distribution using combination of spatially limited photolysis, Ca^{2+} imaging, and patch-clamp recording (protocol), 85–89
materials, 85–86
method, 86–87, 87f
recipes, 88–89
troubleshooting, 88
light sources for, 77
studying activation of epithelial ion channels using global whole-field photolysis (protocol), 79–84
materials, 79–80
method, 80–83, 81f
recipes, 83–84
troubleshooting, 83
in two-photon photolysis combined with kilobeam array scanner to probe calcium signaling in cardiomyocytes (protocol), 116–120
- Ca^{2+} -induced Ca^{2+} release (CICR), 555–556, 559
- Calcineurin, in yeast, 447–450, 449f
- Calcineurin-NFAT signaling cascade, 411
- Calcium (Ca^{2+})
control of intracellular concentration, 93–94
diffusion constant of, 95
as second messenger, 93
- Calcium-ATPases (Ca^{2+} -ATPases)
plasma membrane Ca^{2+} ATPase (PMCA), 231, 277, 291, 293, 438, 446
secretory pathway Ca^{2+} ATPase (SPCA), 231, 446
- SERCA (*see* Sarco-endoplasmic Ca^{2+} ATPases) in yeast, 446
- Calcium (Ca^{2+})-binding kinetics of proteins, 533–551
challenges in studying, 533–534
collecting data to determine the Ca^{2+} -binding properties of DM-nitrophen and proteins (protocol), 546–551
discussion, 549–550
materials, 546
method, 547–549, 547f
recipe, 550
troubleshooting, 549
measuring rate constants of Ca^{2+} indicators (protocol), 541–545
discussion, 545
materials, 541–542, 542t, 543f
method, 543–544
recipe, 545
troubleshooting, 544–545
measuring steady-state properties of Ca^{2+} indicators with a set of calibrated [Ca^{2+}] solutions (protocol), 536–540
discussion, 539–540, 539t
materials, 536
method, 537–538, 537t, 538t
recipe, 540
troubleshooting, 538
overview of method for measuring, 534–535, 534f
- Calcium (Ca^{2+})-binding proteins (CBPs), 533
affinity of, 533
buffering (bCBPs), 533
measuring kinetics of, 533–551
signaling (sCBPs), 533
in yeast, 448–450, 449f
- Calcium blocker cocktail (recipe), 408
- Calcium (Ca^{2+}) channels
bilayer measurement of endoplasmic reticulum, 321–336
black lipid membrane (BLM) experiments to study InsP_3R and RyanR, 322–323
overview, 321–322
preparation of microsomes to study Ca^{2+} channels (protocol), 326–329
harvesting rat cerebella, 327
materials, 326–327
method, 327–328
microsomal preparation, 328
preparation of Sf9 cells, 327–328
recipes, 329
troubleshooting, 328
reconstitution of endoplasmic reticulum InsP_3 receptors into black lipid membranes (protocol), 330–336
materials, 330–331, 331f
method, 331–334, 332f, 334f, 335f
recipes, 336
troubleshooting, 334–335
- Ca^{2+} release-activated Ca^{2+} (CRAC) channels
overview, 277, 279–281, 280f
protocols
measuring endogenous I_{CRAC} and ORAI currents with the patch-clamp technique, 283–290, 287f

- minimum requirements to use calcium imaging to analyze I_{CRAC} , 291–295, 293f
- IP₃ receptors (IP₃Rs), 155–175
- L-type calcium channels (LTCCs), 411–412
- NAADP-targeted (*see* Nicotinic acid adenine dinucleotide phosphate (NAADP))
- patch-clamp electrophysiology of intracellular, 297–319
 - isolating nuclei from cultured cells (protocol), 308–312, 309t, 311f
 - discussion, 311
 - materials, 308–309, 309t
 - method, 309–311, 311f
 - recipe, 312
 - nuclear patch-clamp protocol, 313–318
 - cytoplasmic-side-out configuration, 315–316
 - luminal-side-out configuration, 315
 - materials, 313–314
 - on-nucleus configuration, 314–315
 - patching inner nuclear membrane, 316
 - overview, 297
 - studying single InsP₃R channel activity
 - in inner nuclear membrane, 303–304
 - methods for, 298–300, 299f
 - in outer nuclear membrane, 300–303, 301f, 302f
 - studying single RyR channels in outer nuclear membrane, 304, 304f
 - second-messenger-operated channels (SMOCs), 278
 - store-operated calcium channels (SOCCs), 278, 278f
 - "stretch-activated," 447
 - transient receptor potential (TRP) channels, 277, 278
 - types, 277–278
 - VDAC1 (*see* Voltage-dependent anion-selective channel protein 1)
 - voltage-dependent Ca²⁺ channels (VDCC), 555–556
 - voltage-sensitive
 - overview, 257–262
 - functional diversity of, 259–260, 260t
 - gating transitions of, 262
 - molecular structure of, 260–262, 261f
 - patch-clamp recordings, 257–276
 - whole-cell recordings of Ca²⁺ currents from isolated neonatal dorsal root ganglion (DRG) neurons (protocol), 264–270
 - whole-cell recordings of recombinant channels heterologously expressed in HEK-293 cells (protocol), 271–275, 273f
 - in yeast, 445–451, 446f, 449f, 451f
- Calcium-chelator solution for metal-chelate buffer system (recipe), 72
- Calcium (Ca²⁺) concentration
 - calibration of fluorescence into, 36–37
 - converting fluorescent data into Ca²⁺ concentration (protocol), 46–49
 - intracellular, calcium signaling and, 277–278
 - measuring steady-state properties of Ca²⁺ indicators with a set of calibrated [Ca²⁺] solutions (protocol), 536–540, 537t–539t
 - role in signaling (*see* Calcium (Ca²⁺) signaling)
 - total cellular Ca²⁺ measurements in yeast using flame photometry (protocol), 460–462
- Calcium (Ca²⁺) dynamics
 - analyzing Ca²⁺ dynamics in intact epithelial cells using spatially limited flash photolysis (protocol), 90–92, 91f
 - models of intracellular signaling/dynamics, 553–569
 - compartmentalized models, 555–557
 - exploring oscillations in a point model of intracellular Ca²⁺ concentration (protocol), 562–565, 564f
 - fire-diffuse-fire (FDF) models, 558–559, 559f
 - ordinary differential equation (ODE) models, 554–555, 562–565
 - spatially extended models, 557–559, 559f
 - threshold models, 558
 - time to blip-stochastic simulation of single channel opening (protocol), 566–569, 567f, 568f
 - two-dimensional models, 558–559, 559f
 - whole-cell models, 554–555
 - in plants
 - analysis of, 467–468
 - high-resolution imaging of cytoplasmic Ca²⁺ dynamics in *Arabidopsis* roots (protocol), 474–478
 - imaging of mitochondrial and nuclear Ca²⁺ dynamics in *Arabidopsis* roots (protocol), 484–488
 - live cell imaging of cytoplasmic and nuclear Ca²⁺ dynamics in *Arabidopsis* roots (protocol), 479–483
 - live cell imaging of cytoplasmic Ca²⁺ dynamics in *Arabidopsis* guard cells (protocol), 469–473
 - Calcium (Ca²⁺) fluxes
 - ⁴⁵Ca²⁺ fluxes in intact cells
 - measurement of intracellular Ca²⁺ release (protocol), 225–229
 - technique overview, 216–217
 - ⁴⁵Ca²⁺ fluxes in permeabilized cells
 - analysis of, 215–216, 216f
 - measurement of intracellular Ca²⁺ release (protocol), 219–224
 - technique overview, 213–215
 - mitochondrial Na⁺/Ca²⁺ exchange assays (protocol), 362–366
 - discussion, 364
 - materials, 362–363
 - method, 363–364, 364f
 - recipes, 365–366
 - troubleshooting, 364
 - Calcium-free Ringer's solution for *Xenopus* neurons (recipe), 408
 - Calcium Green, 140, 144
 - Calcium homeostasis, in yeast, 445–448
 - Calcium/hydrogen (Ca²⁺/H⁺) exchanger (antipport), 338, 339
 - Vcx1 protein/gene in yeast, 446f, 447, 448t, 449, 449f
 - Calcium (Ca²⁺) imaging, in plants, 465–488
 - Ca²⁺ dynamics, 467–468
 - high-resolution imaging of cytoplasmic Ca²⁺ dynamics in *Arabidopsis* roots (protocol), 474–478
 - data analysis, 477
 - discussion, 477
 - imaging, 475–477
 - materials, 474–475
 - method, 475–477, 476f
 - plant growth, 475
 - sample preparation, 475, 476f
 - troubleshooting, 477
 - imaging of mitochondrial and nuclear Ca²⁺ dynamics in *Arabidopsis* roots (protocol), 484–488
 - CLSM analysis, 486
 - discussion, 487
 - imaging analyses, 486
 - materials, 484–485
 - method, 485–486
 - recipes, 487
 - sample preparation, 486
 - seed sterilization and growth of plantlets, 485–486
 - troubleshooting, 487
 - live cell imaging of cytoplasmic and nuclear Ca²⁺ dynamics in *Arabidopsis* roots (protocol), 479–483
 - Ca²⁺ imaging, 480
 - data analyses, 482
 - discussion, 482
 - materials, 479–480
 - method, 480–482, 481f
 - plant growth, 480
 - recipes, 483
 - sample preparation, 480, 481f
 - troubleshooting, 482
 - live cell imaging of cytoplasmic Ca²⁺ dynamics in *Arabidopsis* guard cells (protocol), 469–473
 - data analysis, 472
 - discussion, 473
 - imaging, 471–472
 - materials, 469–470
 - method, 470–472, 471f
 - plant growth, 470
 - recipe, 473
 - sample preparation, 470–471, 471f
 - troubleshooting, 472–473
 - overview, 465–486
 - Yellow Cameleon Ca²⁺ indicators, 466–467, 466f
- Calcium (Ca²⁺) indicators
 - analyzing imaging data with ImageJ, 39–40
 - analyze intensity over time, 40
 - background subtraction, 39
 - importing data, 39
 - choosing appropriate, 26–30
 - affinity of indicator for Ca²⁺, 28
 - cell damage and indicator tolerance, 29
 - dual-wavelength Ca²⁺ indicators, 26–27
 - fluorescence intensity, factors influencing, 26
 - imaging hardware, 28–29
 - indicator leakage, 30
 - photobleaching and photodamage, 29–30
 - ratiometric indicators, 26–27
 - single-wavelength Ca²⁺ indicators, 27–28
 - converting fluorescent data into Ca²⁺ concentration (protocol), 46–49
 - concentration determination, 48
 - materials, 46–47
 - obtaining minimum and maximum fluorescence data, 47–48
 - preparation of calibration solution, 47
 - troubleshooting, 48–49
 - data processing and calibration, 33–37
 - background removal, 34, 35, 35f, 36f
 - calibration of fluorescence into Ca²⁺ concentration, 36–37
 - data presentation, 37, 37f
 - image storage, 33–34
 - uneven illumination, 34–35, 36f, 49
 - genetically encoded Ca²⁺ indicators (GECIs), 51–73
 - Yellow Cameleons, 465–466, 479, 482
 - imaging parameters, 30–33
 - bit depth, 31–32
 - detector settings
 - gain, 32–33, 32f
 - offset, 32f, 33
 - look-up table, 33
 - objective lens, 30–31, 31t
 - pixel size, 31
 - loading fluorescent Ca²⁺ indicators into living cells (protocol), 42–45
 - loading cells with ester form of Ca²⁺ indicator, 43–44
 - materials, 42–43
 - setting up imaging experiment, 44

- Calcium (Ca^{2+}) indicators (*Continued*)
troubleshooting, 44–45
for measuring Ca^{2+} -binding kinetics
collecting data to determine the Ca^{2+} -
binding properties of DM-
nitrophen and proteins (protocol),
546–551, 547f
measuring rate constants of Ca^{2+} indicators
(protocol), 541–545, 542t, 543f
measuring steady-state properties of Ca^{2+}
indicators with a set of calibrated
[Ca^{2+}] solutions (protocol),
536–540, 537t–539t
overview, 534–535, 534f
overview, 25–26
sample handling
mounting, 37–38, 38f
perfusion, 38, 38f
solution exchange, 38–39
- Calcium (Ca^{2+}) oscillations, 553, 558, 562–565,
564f
- Calcium oxalate precipitation, 232–233, 245, 252
- Calcium (Ca^{2+}) puffs, 555, 559, 566, 568
- Calcium (Ca^{2+}) pump
back inhibition of, 232
conformations of, 232
mechanisms and features of action, 231–232
phosphorylation assay, 239–240
protocols
high-throughput measurement of Ca^{2+} -
dependent ATPase in COS
microsomes, 234–244
 Ca^{2+} -dependent ATPase-measurement
in 96-well plate format, 236–238,
237f
calculation of ATPase activity, 238, 239f
calculation of turnover rate, 240
cell culture and transfection, 236
discussion, 241–242
materials, 234–236, 235t
method, 236–240
microsome preparation, 236
phosphorylation assay, 239–240
recipes, 242–244
troubleshooting, 240–241
western blotting, 239
measuring Ca^{2+} -dependent Ca^{2+} -uptake
activity in the mouse heart,
245–254
analysis of data, 249–250
 $^{45}\text{Ca}^{2+}$ -uptake measurements,
248–249, 249f
discussion, 252–253
isolation of mouse ventricles, 246
materials, 245–246
method, 246–251
preparation of Ca^{2+} -uptake assay, 247
recipes, 253–254
tissue homogenization, 247–248
troubleshooting, 251–252
western blotting using cardiac
homogenates, 250–251, 250f
strategies for measuring transporter activity,
232–233
types of Ca^{2+} -transport ATPases, 231
- Calcium (Ca^{2+}) signaling
aequorin introduction into zebrafish embryos
for recording, 193–210
calcium-binding proteins (CBPs) and, 533
cyclic adenosine dinucleotide phosphate ribose
(cADPR) and, 491f
glucose-induced, 450–451, 451f
intracellular, 211–212 (*see also* Intracellular
calcium (Ca^{2+}))
stochastic simulation of single channel
opening, 566–569, 567f, 568f
translating into models, 553–569
compartmentalized models, 555–557
exploring oscillations in a point model
of intracellular Ca^{2+} concentration
(protocol), 562–565, 564f
fire-diffuse-fire (FDF) models,
558–559, 559f
ordinary differential equation (ODE)
models, 554–555, 562–565
spatially extended models, 557–559,
559f
threshold models, 558
time to blip-stochastic simulation of
single channel opening (protocol),
566–569, 567f, 568f
two-dimensional models, 558–559,
559f
whole-cell models, 554–555
- investigations using confocal and two-photon
microscopy, 113–138
confocal FLIM of genetically encoded FRET
sensors of quantitative Ca^{2+}
imaging (protocol), 134–138
multibeam two-photon imaging of fast
 Ca^{2+} signals in Langendorff mouse
heart (protocol), 121–125
multichannel imaging of cellular signaling
(protocol), 126–129
overview, 113–114
two-dimensional imaging of fast
intracellular Ca^{2+} release
(protocol), 130–133
two-photon photolysis combined with
kilobeam array scanner to probe
calcium signaling in
cardiomyocytes (protocol),
116–120
- IP_3 receptors (IP_3Rs) and, 155–156
overview, 277–278
as part of integrated signaling network,
412–413
- Xenopus* oocyte as single-cell model for
studying, 381–398
- in yeast, 445–464
calcium homeostasis, 445–448
calcium transport, 446–448, 448t
measuring calcium concentration, 452
measuring calcium uptake using ^{45}Ca
(protocol), 463–464
materials, 463
method, 463–464
monitoring yeast intracellular Ca^{2+} levels
using in vivo bioluminescence assay
(protocol), 456–459
materials, 456–457
method, 457–459, 458f
troubleshooting, 459
signaling pathways, 448–452
calcium-binding proteins, 448–450,
449f
glucose-induced signaling, 450–451,
451f
intracellular calcium modification,
450–452, 451f
programmed cell death, 451–452
total cellular Ca^{2+} measurements using
flame photometry (protocol),
460–462
materials, 460–461
method, 461–462
troubleshooting, 462
- Calcium (Ca^{2+}) sparks, measuring in cardiac
myocytes (protocol), 430–437
analysis of Ca^{2+} -spark recordings, 435
discussion, 434–436
evaluating ryanodine receptor function in intact
cells, 431–432
evaluating ryanodine receptor function in
permeabilized cells, 433–434, 433f
intact *versus* permeabilized cells, 434–435
materials, 430–431
method, 431–434, 433f
recipes, 436–437
SR Ca^{2+} leak, 435
- Calcium transients
in cardiac myocytes, 414, 417–418, 420–421,
422f, 438, 441–442
in developing *Xenopus* spinal neurons
(protocol), 399–409
discussion, 407–408
materials, 399–400
method, 400–407
data analysis, 404
general imaging methodology,
401–402, 402f
imaging filopodial transients in vivo,
403–404, 403f
imaging growth cone transients in vivo,
402–403, 403f
imposition of Ca^{2+} spikes, 405, 405f
manipulation of Ca^{2+} dynamics in
growth cones, 406
manipulation of filopodial Ca^{2+}
dynamics, 406–407
perturbation experiments, 404
preparation of dissociated cell cultures,
400–401
preparation of spinal cords, 401
recipes, 408–409
troubleshooting, 407
- Calcium (Ca^{2+}) transport
channels (*see* Calcium (Ca^{2+}) channels)
mitochondrial (*see* Mitochondrial Ca^{2+}
transport)
in yeast, 446–448, 448t
- Calcium (Ca^{2+}) uptake
measuring in yeast using ^{45}Ca (protocol),
463–464
- Calcium (Ca^{2+}) wave
intercellular (*see* Intercellular Ca^{2+} wave
(ICW))
intracellular, 553–554, 557–559, 559f
- Calmodulin, 222
 Ca^{2+} -binding kinetics, 534
cyclic adenosine dinucleotide phosphate ribose
(cADPR)-mediated Ca^{2+} release
and, 490
in yeast, 446, 448–450, 449f, 452
Yellow Cameleon indicators and, 466, 467
- Calnexin, 211
- Ca^{2+} loading solution for measuring Ca^{2+} fluxes
across the mitochondrial inner
membrane (recipe), 365
- Calreticulin, 211
- Cameleons. *See also* Yellow Cameleon indicators
binding parameters of, 52t
use in yeast, 452
- Camera
complementary metal oxide semiconductor
(CMOS), 20, 28, 436
electron-multiplying charge coupled device
(EMCCD), 13, 20, 28, 111, 436
- Ca^{2+} mix (recipe), 242
- CaMKII, 411
- Capacitative Ca^{2+} influx, measurement of, 188–189
- Carbachol, 188
- Carbonyl cyanide *m*-chlorophenyl hydrazone
(CCCP), 291, 293
- Carboxymethyl (CM)-cellulose, voltage-dependent
anion-selective channel protein 1
(VDAC1) purification using,
345–346
- Cardiomyocytes (cardiac myocytes)
compartmentalized models of Ca^{2+} signaling
and, 555, 556
excitation-contraction coupling, 411–412, 412f
systematic approach for assessing Ca^{2+} handling
in, 411–443

- assessing Ca^{2+} -removal pathways in cardiac myocytes (protocol), 438–443
 - discussion, 441–442
 - materials, 438–439
 - method, 439–441, 440f, 441f
 - recipes, 442–443
- characterizing trigger for sarcoplasmic reticulum Ca^{2+} release in cardiac myocytes (protocol), 420–424, 422f
 - discussion, 421–423, 422f
 - materials, 420
 - method, 421
 - recipes, 423–424
- measuring Ca^{2+} sparks in cardiac myocytes (protocol), 430–437
 - analysis of Ca^{2+} -spark recordings, 435
 - discussion, 434–436
 - evaluating ryanodine receptor function in intact cells, 431–432
 - evaluating ryanodine receptor function in permeabilized cells, 433–434, 433f
 - intact *versus* permeabilized cells, 434–435
 - materials, 430–431
 - method, 431–434, 433f
 - recipes, 436–437
 - SR Ca^{2+} leak, 435
- measuring sarcoplasmic reticulum Ca^{2+} content, fractional release, and Ca^{2+} buffering in cardiac myocytes (protocol), 425–429, 427f
 - discussion, 426–428, 427f
 - materials, 425
 - method, 426
 - recipes, 428–429
 - troubleshooting, 426
- monitoring intracellular Ca^{2+} in cardiac myocytes using Fluo-3 (protocol), 414–419, 415f
 - discussion, 417–418
 - materials, 414–415, 415f
 - method, 416–417
 - recipes, 418
 - troubleshooting, 417
- two-dimensional imaging of fast intracellular Ca^{2+} release in (protocol), 130–133, 132f
- two-photon photolysis combined with kilobeam array scanner to probe calcium signaling in (protocol), 116–120, 118f
- Ca^{2+} release-activated Ca^{2+} (CRAC) channels
 - overview, 277, 279–281, 280f
 - protocols
 - measuring endogenous I_{CRAC} and ORAI currents with the patch-clamp technique, 283–290, 287f
 - minimum requirements to use calcium imaging to analyze I_{CRAC} , 291–295, 293f
- Ca^{2+} signatures, 465
- Ca^{2+} uptake solution for assaying Ca^{2+} transport in mitochondria (recipe), 361
- CBPs. *See* Calcium (Ca^{2+})-binding proteins
- Cch1 channels, 445, 446, 446f, 447, 448t, 449f
- CD38, 489–490, 491f
 - measuring CD38 activity by reverse-phase HPLC (protocol), 498–502
- Cell wall integrity (CWI) Pkc1/Slt2 pathway, 450
- Central core disease, 323
- Cerebella, harvesting rat, 327
- CFP, in Yellow Cameleons, 466, 466f, 471–472, 471f, 475, 477, 480, 482, 486–487
- Channel conductance, recording through voltage-dependent anion-selective channel protein 1 (VDAC1), 349–354, 351f, 352f
- Channelopathies, 447
- Channels. *See* Calcium (Ca^{2+}) channels
- Charcoal, 496
- Chelator solution for metal-chelate buffer system (recipe), 72
- Chelex-treated ultrapure H_2O (recipe), 63, 72
- Chemicals, properties of common hazardous, 575–576
- Chimeras, aequorin, 181t–182t
- Chromatic aberration, 7
- CICR (Ca^{2+} -induced Ca^{2+} release), 555–556, 559
- Ci-InsP₃, 90–91
- Cis recording solution (recipe), 336
- Coelenterates, 177
- Coelenterazine, 178–179, 178f, 180, 187–190, 193–194, 203, 209
 - loading yeast with, 452, 457, 459
- Collagenase B, 402
- Complementary metal oxide semiconductor (CMOS) camera, 20, 28, 436
- Concentration. *See* Calcium (Ca^{2+}) concentration
- Confocal laser scanning microscopy (CLSM)
 - basic elements of, 8f
 - complications of, 9
 - dwell time, 9
 - image speed, 10–11, 10f
 - imaging Ca^{2+} dynamics in plants
 - high-resolution imaging of cytoplasmic Ca^{2+} dynamics in *Arabidopsis* roots (protocol), 474–478, 476f
 - imaging of mitochondrial and nuclear Ca^{2+} dynamics in *Arabidopsis* roots (protocol), 484–488, 485f
 - live cell imaging of cytoplasmic and nuclear Ca^{2+} dynamics in *Arabidopsis* roots (protocol), 479–483, 481f
 - lasers, 12
 - line scan, 11
 - measuring Ca^{2+} sparks in cardiac myocytes (protocol), 430–437
 - photomultiplier gain, 11–12
 - pinhole size, 9–10, 9f
 - principles of operation, 7–8, 8f
 - spatially limited photolysis, 77
 - spectral imaging, 12–13
- Confocal microscopy
 - calcium signaling investigations using, 113–138
 - confocal FLIM of genetically encoded FRET sensors of quantitative Ca^{2+} imaging (protocol), 134–138
 - discussion, 137–138
 - materials, 134–135
 - method, 135–137, 136f
 - recipe, 138
 - troubleshooting, 137
 - multibeam two-photon imaging of fast Ca^{2+} signals in Langendorff mouse heart (protocol), 121–125
 - discussion, 124
 - materials, 121–122
 - methods, 122–124, 123f
 - recipes, 124
 - troubleshooting, 124
 - multichannel imaging of cellular signaling (protocol), 126–129
 - discussion, 129
 - materials, 126–127
 - method, 127–128, 128f
 - recipe, 129
 - troubleshooting, 128–129
 - overview, 113–114
 - two-dimensional imaging of fast intracellular Ca^{2+} release (protocol), 130–133
 - discussion, 131, 133
 - materials, 130
 - method, 131, 132f
 - recipe, 133
- troubleshooting, 131
- two-photon photolysis combined with kilobeam array scanner to probe calcium signaling in cardiomyocytes (protocol), 116–120
 - discussion, 119
 - materials, 116–117
 - method, 117–118, 118f
 - recipes, 119–120
 - troubleshooting, 118–119
- optical sectioning, 29
- overview, 7–13
 - basic elements of, 8f
 - complications of, 9
 - dwell time, 9
 - goal of, 8
 - image speed, 10–11, 10f
 - lasers, 12
 - line scan, 11
 - photomultiplier gain, 11–12
 - pinhole size, 9–10, 9f
 - principles of operation, 7–8, 8f
 - spectral imaging, 12–13
 - spinning disc confocal microscopy, 13
 - pixel size, 31
- Connexin 43 hemichannels, 490
- Connexin (Cx) proteins, 94–96, 99–101
- CoroNa-Red, 363–364, 364t
- Coupling gain, 427–428
- Coverslips
 - diameter and size of, 37
 - transferring to imaging chamber, 38
- CPA, 292
- cpVenus, 486–487
- Creatine kinase, 505
- Creatine phosphokinase, 513, 522
- Crz1, 449
- Cs-aspartate-based internal pipette solution (recipe), 423, 429
- CsCl-based internal pipette solution (recipe), 442
- Cs-Tyrode-based external solution (recipe), 423, 429, 443
- Ctenophores, 177
- Culture medium for cells expressing recombinant Ca^{2+} channels (recipe), 275
- Culture medium for DRG neurons (recipe), 269
- Currents
 - I_{CRAC} (Ca^{2+} release-activated Ca^{2+} current), 277–295
 - activation of, 278–279
 - measuring endogenous I_{CRAC} and ORAI currents with the patch-clamp technique (protocol), 283–290, 287f
 - minimum requirements to use calcium imaging to analyze I_{CRAC} (protocol), 291–295, 293f
 - K^+ , 427
 - L-type Ca^{2+} current (I_{CaL}), 420–423
 - Na^+ , 423
 - $\text{Na}^+/\text{Ca}^{2+}$ exchanger current (I_{NCX}), 425–428, 427f, 439–441, 440f, 441f
 - recording through voltage-dependent anion-selective channel protein 1 (VDAC1), 349–354, 351f, 352f
- Cutting devices, safe handling of, 573
- Cyclic adenosine dinucleotide phosphate ribose (cADPR), 489–507, 509, 510
 - caged, 528
 - Ca^{2+} release evoked by
 - measuring, 503–507
 - release from sea urchin egg homogenates, 514f
 - Ca^{2+} release from sea urchin egg homogenates, 514f
 - cycling assay determining intracellular levels (protocol), 493–498
 - discussion, 496

- Cyclic adenosine dinucleotide phosphate ribose (cADPR), (*Continued*)
 materials, 493–494
 method, 494–495
 for adherent cells, 494–495
 for cells in suspension, 494
 troubleshooting, 495–496
 degradation of, 490
 hydrolase activity, measuring, 498–500
 measuring Ca^{2+} release evoked by (protocol), 503–507
 discussion, 506
 materials, 503–504
 method, 504–506
 measuring in permeabilized cells, 504–505
 microinjection of cADPR and Ca^{2+} imaging, 505–506
 recipes, 507
 troubleshooting, 506
 measuring CD38 activity by reverse-phase HPLC (protocol), 498–502
 discussion, 500–501
 materials, 498–499
 method, 499–500
 recipes, 501
 troubleshooting, 500
 metabolism of, 490
 overview, 489
 structure of, 490f
 topological paradox, 490
- Cycling assay determining intracellular cyclic ADP-ribose levels (protocol), 493–498
- Cyclopiazonic acid, 161, 188–189, 241
- Cyp D, 338
- Cytochrome C, 99
- Cytosolic Ca^{2+} , measurement of, 187–188
- D**
- D1, binding parameters of, 62t
- Danieau's solution (recipe), 204, 210
- Darier disease, 241
- D3cpV, binding parameters of, 62t
- Dechorionating embryos, 200
- Department of Health, Education, and Welfare (HEW), U.S., 575
- Dextran and leakage-resistant Ca^{2+} indicators, 30
- De Young-Keizer model, 562–565
- D-family cameleons, binding parameters of, 52t
- Diaclylglycerol (DAG), 94, 450, 451f
- Dialysis buffer for purification of VDACL1 from mitochondria (recipe), 348
- Diaphorase, 495
- 1(2-nitrophenyl)diazoethane, 526
- DibromoBAPTA, affinity for Ca^{2+} , 537t
- Dichroic mirror, 4, 5
- DIDS, 421, 426, 427
- Diffraction limit, 3
- Diffusion coefficient, 557
- Digitonin, 58, 63, 68, 188–190, 213, 364
- Dihydropyridines, 259, 260t
- Diisopropylethylamine (DIEA), 518–519
- DinitroBAPTA, affinity for Ca^{2+} , 537t
- Disposal
 general cautions, 571–573
 of laboratory waste, 573
- Dissociating medium for DRG neurons (recipe), 269
- Dissociation constant (K_d) of indicator for Ca^{2+} , 28
 measuring the in situ K_d of GECl (protocol), 65–73
- 2,5-di-*t*-butyl-1,4-benzohydroquinone, 241
- Divalent-cation-free minimal medium (recipe), 408
- DM-nitrophen
 affinity of, 550
 collecting data to determine the Ca^{2+} -binding properties of DM-nitrophen and proteins (protocol), 546–551, 547f
- measuring rate constants of Ca^{2+} indicators (protocol), 541–545, 542t, 543f
- preparation of stock, 543
- in two-photon photolysis combined with kilobeam array scanner to probe calcium signaling in cardiomyocytes (protocol), 116–119
- DMSO
 sulfinpyrazone, 45
 use in electroporation loading, 106
- Dorsal root ganglion (DRG) neurons
 dissecting, 265, 266f
 L-type Ca^{2+} current in, 260
 N-type Ca^{2+} current in, 260
 whole-cell patch-clamp recordings of Ca^{2+} currents from isolated neonatal dorsal root ganglion (DRG) neurons (protocol), 264–270
- Dounce homogenizer, 311, 513
- DsRed in multichannel imaging of cellular signaling (protocol), 126–129, 128f
- Dual emission detectors, 27
- Duall homogenizer, 311
- Dual-wavelength Ca^{2+} indicators, 26–27
- Dwell time, in confocal microscopy, 9
- Dyes, hazards of, 575
- E**
- EDTA, affinity for Ca^{2+} , 537t
- Efflux medium for intact cells (recipe), 228
- Efflux medium for permeabilized cells (recipe), 222
- EGTA
 affinity for Ca^{2+} , 537t
 in Ca^{2+} sparks measurement, 435
 Ca^{2+} store depletion by, 279, 286
- EGTA (0.5 M) (recipe), 63, 72
- EGTA Ca^{2+} - free artificial seawater (recipe), 515
- EGTA mix (recipe), 242
- EGTA solution (100 mM) (recipe), 436
- Electrodes, Ca^{2+} -sensitive, 367–380
 advantages of, 369, 370
 disadvantages of, 370
 history of, 369
 making microelectrodes (protocol), 376–380
 discussion, 379–380, 380f
 double-barreled electrodes, 378–379
 materials, 376–377
 single-barreled electrodes, 377–378
 troubleshooting, 379
 making minielectrodes (protocol), 372–375
 discussion, 374–375
 materials, 372–373
 method, 373–374, 374f
 troubleshooting, 374
 PVC use in, 369–370
 response time and Ca^{2+} indicator choice, 368–369
 uses of, 370
 in vitro calibration curves, 368f
- Electron-multiplying charge coupled device (EMCCD), 13, 20, 38, 111, 436
- Electroporation loading
 bipolar pulsed electroporation, 97
 electrical pulse protocol, 97
 loading efficiency, 97, 106, 107
 of membrane-impermeable molecules (protocol), 104–108
 materials, 104–105
 method, 105–106
 electroporation, 105–106
 loading efficiency, 106
 recipes, 107
 troubleshooting, 106–107
 overview, 97–101
 setup, 97, 98f
- single-cell electroporation, 97
- Electroporation of C6 glioma cells, 99t
- Embryos
 aequorin introduction into zebrafish, 193–210, 195f
 microinjecting holo-aequorin into dechorionated and intact zebrafish embryos (protocol), 197–205
 reconstitution of holo-aequorin with apoaequorin mRNA and coelenterazine in zebrafish embryos (protocol), 206–210
 dechorionating, 200
- Emission filter, 3f, 4
- ENAI gene, 450
- Endolysosomal channels, NAADP and, 509, 510
- Endoplasmic reticulum, measurement of Ca^{2+} in, 190
- Endoplasmic reticulum Ca^{2+} channels, 321–336
 black lipid membrane (BLM) experiments to study InsP₃R and RyanR, 322–323
 overview, 321–322
 preparation of microsomes to study Ca^{2+} channels (protocol), 326–329
 harvesting rat cerebella, 327
 materials, 326–327
 method, 327–328
 microsomal preparation, 328
 preparation of Sf9 cells, 327–328
 recipes, 329
 troubleshooting, 328
 reconstitution of endoplasmic reticulum InsP₃ receptors into black lipid membranes (protocol), 330–336
 materials, 330–331, 331f
 method, 331–334, 332f, 334f, 335f
 recipes, 336
 troubleshooting, 334–335
- Environmental Protection Agency (EPA), U.S., 573
- EPSPs (excitatory postsynaptic potentials), 142–143, 142f
- ETH 129 sensor, 369, 373, 377
- ETH 1001 sensor, 369
- European *Saccharomyces cerevisiae* Archives of Functional Analysis (EUROSCARF), 445
- EUROSCARF (European *Saccharomyces cerevisiae* Archives of Functional Analysis), 445
- Evanescence wave, 17, 20, 28
- Exchanges of media/solutions, 38–39
- Excitation-contraction coupling, 411–412, 412f
- Excitation filter, 3f, 4
- Excitation wavelength selection, in two-photon microscopy, 15
- Excitatory postsynaptic potentials (EPSPs), 142–143, 142f
- External recording solution for Ca^{2+} currents (recipe), 269
- External recording solution for recombinant Ca^{2+} channels (recipe), 275
- Extracellular buffer (recipe), 501
- Extracellular solution with 1 mM Ca^{2+} (recipe), 294
- Extracellular solution with no free Ca^{2+} (recipe), 295
- F**
- Fatty acid oxidation, 338
- Fibronectin, 401
- Fig1 protein/gene, 446f, 447, 448t, 449f
- Filter cube, 4
- Filters
 blocking, 4–5
 light separation and, 3, 3f
 orientation of, 5
 polarizing to attenuate laser power, 12
 sources of, 3

- FITC-IP₃, 173–174, 173f, 174f
 Fks2 protein/gene, 450
 Flame photometry, total cellular Ca²⁺
 measurements in yeast using
 (protocol), 460–462
 materials, 460–461
 method, 461–462
 troubleshooting, 462
 Flash photolysis of caged IP₃ to trigger intercellular
 Ca²⁺ waves (protocol), 109–112
 materials, 109–110
 methods, 110–111
 ester loading of fluorescent Ca²⁺ indicator,
 110
 quantification of ICW size, 111
 uncaging of IP₃ using flash photolysis,
 110–111
 recipe, 112
 troubleshooting, 111–112
 FlexStation 3, 163, 164
 FLIM (fluorescence lifetime imaging), confocal of
 genetically encoded FRET sensors
 of quantitative Ca²⁺ imaging
 (protocol), 134–138
 discussion, 137–138
 materials, 134–135
 method, 135–137, 136f
 recipe, 138
 troubleshooting, 137
 Fluo-3
 Fluo-3-AM
 calcium transients in developing *Xenopus*
 spinal neurons (protocol),
 399–409, 403f
 in flash photolysis of caged IP₃ to trigger
 intercellular Ca²⁺ waves (protocol),
 109–111
 monitoring intracellular Ca²⁺ in cardiac
 myocytes using Fluo-3 (protocol),
 414–419
 measuring Ca²⁺ release from sea urchin egg
 homogenates, 514
 measuring Ca²⁺ sparks in cardiac myocytes
 (protocol), 432–434
 peak excitation and emission wavelengths, 417
 Fluo-4
 accumulation in mitochondria, 30
 Fluo-4-AM
 in analyzing Ca²⁺ dynamics in intact
 epithelial cells using spatially
 limited flash photolysis (protocol)
 materials, 90–91
 calcium transients in developing *Xenopus*
 spinal neurons (protocol),
 399–409, 403f
 in multibeam two-photon imaging of fast
 Ca²⁺ signals in Langendorff mouse
 heart (protocol), 122
 in multichannel imaging of cellular
 signaling (protocol), 126–129, 128f
 in two-dimensional imaging of fast
 intracellular Ca²⁺ release
 (protocol), 130–133, 132f
 Fluo-4FF, 539
 as single-wavelength Ca²⁺ indicators, 27–28
 in two-photon photolysis combined with
 kilobeam array scanner to probe
 calcium signaling in
 cardiomyocytes (protocol),
 116–119, 118f
 variations in batches of, 539
 Fluo-8-AM, in multibeam two-photon imaging of
 fast Ca²⁺ signals in Langendorff
 mouse heart (protocol), 122
 Fluo-3-AM solution (recipe), 409
 Fluo-4-AM solution (recipe), 409
 Fluorescence, principles of, 2–3, 2f
 Fluorescence intensity, factors influencing, 26
 Fluorescence lifetime imaging (FLIM), confocal of
 genetically encoded FRET sensors
 of quantitative Ca²⁺ imaging
 (protocol), 134–138
 discussion, 137–138
 materials, 134–135
 method, 135–137, 136f
 recipe, 138
 troubleshooting, 137
 Fluorescence light separation, 3–5
 blocking filters, 4–5
 dichroic mirror, 4
 filter cube, 4
 multiple fluorophores, 5
 orientation of dichroic mirrors and filters, 5
 Fluorescence microscopy, 1–23
 confocal microscopy
 overview, 7–13
 basic elements of, 8f
 complications of, 9
 dwell time, 9
 goal of, 8
 image speed, 10–11, 10f
 lasers, 12
 line scan, 11
 photomultiplier gain, 11–12
 pinhole size, 9–10, 9f
 principles of operation, 7–8, 8f
 spectral imaging, 12–13
 spinning disc confocal microscopy, 13
 fluorescence light separation, 3–5
 blocking filters, 4–5
 dichroic mirror, 4
 filter cube, 4
 multiple fluorophores, 5
 orientation of dichroic mirrors and filters, 5
 light paths, basic, 3, 3f
 principles of fluorescence, 2–3, 2f
 super-resolution microscopy
 overview, 21–23
 PALM/STORM super-resolution
 imaging, 21–23
 STED super-resolution imaging, 23, 23f
 total internal reflection fluorescence (TIRF)
 microscopy
 overview, 17–21
 imaging, 20–21
 prism-based, 18
 safety issues, 20
 sample preparation, 21
 through-the-objective, 18–19, 19f
 two-photon microscopy
 overview, 13–17
 auto-fluorescence, 16
 basic elements of, 14f
 excitation wavelength selection, 15
 implications for design, 14
 laser damage, 14
 photobleaching, 15
 second-harmonic imaging, 16
 trans-illumination imaging, 16–17, 17f
 wide-field
 combined transmitted and fluorescence
 microscopy, 7
 overview, 5–7
 Fluorescent indicators. *See* Calcium (Ca²⁺)
 indicators; *specific indicators*
 Fluorescent proteins
 in high-throughput fluorescence polarization
 assay of ligand binding to IP₃
 receptors protocol, 167–175, 169f
 multiple excitation wavelengths, 15
 PA-FPs, 22
 Fluorophores, multiple, 5
 Förster resonance energy transfer (FRET)
 confocal FLIM of genetically encoded FRET
 sensors of quantitative Ca²⁺
 imaging (protocol), 134–138
 genetically encoded Ca²⁺ indicators (GECIs),
 51–73
 converting fluorescent data into Ca²⁺
 concentration (protocol), 46–49
 Yellow Chameleons, 466–467, 466f, 472, 477
 FRET. *See* Förster resonance energy transfer
 Fura-2
 dissociation constant of, 141
 as dual-wavelength indicator, 26–27
 Fura-2-AM
 calcium imaging to analyze I_{CRAC}, 292, 294
 calcium transients in developing *Xenopus*
 spinal neurons (protocol),
 399–409
 microinjection of cADPR and Ca²⁺
 imaging, 505–506
 isosbestic point, 27
 measuring cADPR-evoked Ca²⁺ release in
 permeabilized cells, 505
 Mn²⁺ binding to, 27
 in yeast, 452
 Fura dyes
 in combining Ca²⁺ and membrane potential
 imaging, 140–143, 141f, 146–149,
 147f
 in combining Ca²⁺ imaging with
 neurotransmitter photorelease, 144
 dissociation constants of, 141
 Fura-FF, 141, 142f, 143
 Fura Red
 advantage of, 27
 Fura-Red-AM in calcium transients in
 developing *Xenopus* spinal neurons
 (protocol), 399–409
- ## G
- Gadolinium, 447
 Gain setting, 32–33, 32f
 Gap junctions, 94–96, 94f
 Gas containers, safe handling of, 572
 GCI promoter, 469
 GECIs. *See* Genetically encoded Ca²⁺ indicators
 Gene expression
 copy numbers of expressed proteins, 383
 Xenopus oocyte, 381–383, 383t, 388–393
 Genetically encoded Ca²⁺ indicators (GECIs),
 51–73
 intensiometric, 51
 measuring the in situ K_d of GECI (protocol),
 65–73
 discussion, 71
 materials, 65–66
 method, 66–70, 67t–68t, 69f–70f
 recipes, 72–73
 troubleshooting, 70–71
 ratiometric
 binding parameters, 53t
 design and use, 51–54, 52f
 verifying function and localization of GECIs
 and converting FRET ratios to Ca²⁺
 concentrations (protocol), 55–64
 materials, 55–56
 method, 56–62, 57f, 60f–61f
 recipes, 63–64
 troubleshooting, 62–63
 Yellow Chameleons, 465–466, 479, 482
 GIC (glucose-induced calcium) transporter, 446f,
 447, 449f
 Glass-bottom imaging dishes, construction of, 56,
 57f
 Glucose-induced calcium (Ca²⁺) signaling,
 450–451, 451f, 457, 458f
 Glutamate, 476
 Glutamate receptors, activating with L-glutamate
 photorelease, 140, 150–152, 151f
 Glutathione, 222
 G protein-coupled receptors (GPCRs), 94f, 95

- Gramicidin, 280, 418
 Group velocity dispersion, 14
 Growth medium (*Arabidopsis*) (recipe), 487
 Guard cells, live cell imaging of cytoplasmic Ca^{2+} dynamics in *Arabidopsis* (protocol), 469–473
- ## H
- HACS (high-affinity calcium system), 446f, 447, 450
 Hailey-Hailey disease, 241
 Halistaurin, 177
 Halothane, 323
 Hanks balanced salt solution-HEPES (recipe), 107, 112
 Hazardous chemicals, properties of common, 575–576
- Heart
 excision and cannulation of mouse heart, 122
 excitation-contraction coupling, 411–412, 412f
 measuring Ca^{2+} -dependent Ca^{2+} -uptake activity in the mouse heart (protocol), 245–254
 multibeam two-photon imaging of fast Ca^{2+} signals in Langendorff mouse heart (protocol), 121–125, 123f
 two-dimensional imaging of fast intracellular Ca^{2+} release (protocol), 130–133, 132f
 two-photon photolysis combined with kilobeam array scanner to probe calcium signaling in cardiomyocytes (protocol), 116–120, 118f
- HEDTA, affinity for Ca^{2+} , 537t
 HEK-293 cells, whole-cell patch-clamp recordings of recombinant voltage-sensitive Ca^{2+} channels heterologously expressed in (protocol), 271–275, 273f
- Heparin, 246
 HEPES-buffered Hanks balanced salt solution (HHBSS) with Ca^{2+} (recipe), 64, 72
 HEPES-buffered Hanks balanced salt solution without Ca^{2+} (Ca^{2+} -free HHBSS) (recipe), 64
 HEPES-buffered Hanks balanced salt solution without Ca^{2+} and Mg^{2+} ($\text{Ca}^{2+}/\text{Mg}^{2+}$ -free HHBSS) (recipe), 73
 HEPES-buffered saline (HBS) (recipe), 165
 Heterologous expression in *Xenopus* oocyte, 381–383, 383t, 388–393
 High-salt buffer (HSB) for microsomal preparation (recipe), 329
 High-throughput screens, history of, 156–157
 Histamine, 188, 190
 Homogenization
 of mouse heart, 247–248
 for nuclei isolation, 310, 311
 of rat liver, 344
 of sea urchin eggs, 513
 Homogenization buffer for Ca^{2+} -uptake assay (recipe), 253
 Homogenization buffer (HB) for cerebellar microsomal preparation (recipe), 329
 Homogenization buffer for sea urchin homogenates (recipe), 515
 Homogenization buffer (SF9HB) for Sf9 microsomal preparation (recipe), 329
- HPLC, measuring CD38 activity by reverse-phase (protocol), 498–502
 discussion, 500–501
 materials, 498–499
 method, 499–500
 recipes, 501
 troubleshooting, 500
- HPLC buffer A (recipe), 501
 HPLC buffer B (recipe), 501
 Hsl1, 450
 Hydrogen/potassium (H^+/K^+) exchanger, 338, 339
 Hydroxyapatite:celite column, for VDACL1 purification, 344–345
 Hypotonic shock in yeast, 445, 450, 457, 458f
- ## I
- IBlot, 398
 I_{CRAC} (Ca^{2+} release-activated Ca^{2+} current), 277–295
 activation of, 278–279
 protocols
 measuring endogenous I_{CRAC} and ORAI currents with the patch-clamp technique, 283–290, 287f
 discussion, 288
 materials, 283–284
 method, 284–287, 287f
 recipes, 289–290
 troubleshooting, 287–288
 minimum requirements to use calcium imaging to analyze I_{CRAC} , 291–295, 293f
 discussion, 294
 materials, 291–292
 method, 292–294, 293f
 recipes, 294–295
 troubleshooting, 294
 I_{CRAC} Ca^{2+} -containing bath solution (recipe), 289
 I_{CRAC} Ca^{2+} -free bath solution (recipe), 289
 I_{CRAC} divalent-free (DVF) bath solution (recipe), 289
 I_{CRAC} pipette solution for active depletion (recipe), 289
 I_{CRAC} pipette solution for passive depletion (recipe), 289
- ICW. See Intercellular Ca^{2+} wave
 IGOR Pro (software), 137
 Illumination
 selective, 28–29
 trans-illumination imaging, 16–17, 17f
 for uncaging IP_3 , 111
 uneven illumination, 34–35, 36f, 49
- ImageJ, 39–40, 435
 analyze intensity over time, 40
 background subtraction, 39
 importing data, 39
 Ratio Plus plug in, 486
- Image speed, in confocal microscopy, 10–11, 10f
 Image storage, 33–34
- Imaging
 analyzing imaging data with ImageJ, 39–40
 analyze intensity over time, 40
 background subtraction, 39
 importing data, 39
 calcium (Ca^{2+}) imaging in plants, 465–488
 Ca^{2+} dynamics, 467–468
 high-resolution imaging of cytoplasmic Ca^{2+} dynamics in *Arabidopsis* roots (protocol), 474–478
 data analysis, 477
 discussion, 477
 imaging, 475–477
 materials, 474–475
 method, 475–477, 476f
 plant growth, 475
 sample preparation, 475, 476f
 troubleshooting, 477
 imaging of mitochondrial and nuclear Ca^{2+} dynamics in *Arabidopsis* roots (protocol), 484–488
 CLSM analysis, 486
 discussion, 487
 imaging analyses, 486
- materials, 484–485
 method, 485–486
 recipes, 487
 sample preparation, 486
 seed sterilization and growth of plantlets, 485–486
 troubleshooting, 487
- live cell imaging of cytoplasmic and nuclear Ca^{2+} dynamics in *Arabidopsis* roots (protocol), 479–483
 Ca^{2+} imaging, 480
 data analyses, 482
 discussion, 482
 materials, 479–480
 method, 480–482, 481f
 plant growth, 480
 recipes, 483
 sample preparation, 480, 481f
 troubleshooting, 482
- live cell imaging of cytoplasmic Ca^{2+} dynamics in *Arabidopsis* guard cells (protocol), 469–473
 data analysis, 472
 discussion, 473
 imaging, 471–472
 materials, 469–470
 method, 470–472, 471f
 plant growth, 470
 recipe, 473
 sample preparation, 470–471, 471f
 troubleshooting, 472–473
 overview, 465–486
 Yellow Cameleon Ca^{2+} indicators, 466–467, 466f
- calcium transients in developing *Xenopus* spinal neurons (protocol), 399–409
- combining calcium imaging with other optical techniques, 139–153
 applications
 combining Ca^{2+} and V_{M} optical measurements, 141–143, 142f
 combining Ca^{2+} imaging with neurotransmitter photorelease, 143–144, 144f
 overview, 139–140, 141f
 protocols
 combining Ca^{2+} and membrane potential imaging in single neurons, 146–149, 147f
 combining Ca^{2+} imaging with L-glutamate photorelease, 150–153, 151f
- data processing and calibration, 33–37
 background removal, 34, 35, 35f, 36f
 calibration of fluorescence into Ca^{2+} concentration, 36–37
 data presentation, 37, 37f
 image storage, 33–34
 uneven illumination, 34–35, 36f, 49
- imaging parameters, 30–33
 bit depth, 31–32
 detector settings
 gain, 32–33, 32f
 offset, 32f, 33
 look-up table, 33
 objective lens, 30–31, 31t
 pixel size, 31
- microinjection of cADPR and Ca^{2+} imaging, 505–506
- minimum requirements to use calcium imaging to analyze I_{CRAC} , 291–295, 293f
- sample handling
 mounting, 37–38, 38f
 perfusion, 38, 38f
 solution exchange, 38–39
- Imaging chamber, 38
 Imaging hardware, 28–29
 Imaging medium (*Arabidopsis*) (recipe), 487

- Imaging solution (recipe), 92
 Immobilon-P membrane plate, 495
 Incubation buffer (recipe), 473
 I_{NCX} (Na^+ / Ca^{2+} exchanger current), 425–428, 427f, 439–441, 440f, 441f
 Indicator leakage, 30
 Indicators. *See also* Calcium (Ca^{2+}) indicators; *specific agents*
 multiple, 5
 spectra of excitation and emission, 3
 Indo-1, 452
 disadvantage of, 27
 dual emission detectors and, 27
 Inositol 1,4,5-trisphosphate (IP_3), 173–174, 173f, 174f
 caged IP_3 , 79, 81f, 90–92, 96, 99–101, 100f, 109–112, 528
 calcium release in yeast, 445, 450–451, 451f
 Ca^{2+} release from sea urchin egg homogenates, 514f
 Ca^{2+} store depletion by, 278, 286
 diffusion of, 95
 gap junction permeability of, 95
 intercellular Ca^{2+} waves and, 94–95, 94f, 96, 109–112
 intracellular Ca^{2+} signaling and, 212, 213
 Inositol 1,4,5-trisphosphate receptor (IP_3R), 99, 155–175, 277, 297
 activation by means of G-protein-coupled receptors, 321–323
 black lipid membrane (BLM) experiments to study IP_3R and RyanR, 322–323
 Ca^{2+} signaling via, 155–156, 212, 213
 channels, 94, 99
 in endoplasmic reticulum, 321–323
 high-throughput fluorescence polarization assay of ligand binding to IP_3 receptors (protocol), 167–175, 169f
 discussion, 173–174, 173f, 174t
 materials, 167–168
 method, 168–172
 anisotropy measurements, 170–171
 data analysis, 171–172
 optimization of FP assay, 168
 preparation of plates for FP equilibrium competition binding assays, 170
 preparation of plates for FP saturation binding assays, 168–170
 recipe, 174
 troubleshooting, 172–173
 high-throughput functional assays of IP_3 -evoked Ca^{2+} release (protocol), 159–166, 162f
 discussion, 164–165
 materials, 159–160
 method, 160–163
 cell culture, 160
 data analysis, 163
 loading cells with Mag-Fluo-4, 161
 measurement of Ca^{2+} uptake and release, 163
 permeabilization of cells, 161, 163
 preparation of assay plates and compound plates, 160–161
 recipes, 165
 troubleshooting, 163–164
 impact of buffers on, 557
 in models of intracellular calcium signaling, 554–560, 566–568, 567f, 568f
 stochastic simulation of single channel opening, 566–568, 567f, 568f
 patch-clamp electrophysiology of channels
 isolating nuclei from cultured cells (protocol), 308–312, 309t, 311f
 nuclear patch-clamp protocol, 313–318
 cytoplasmic-side-out configuration, 315–316
 luminal-side-out configuration, 315
 materials, 313–314
 on-nucleus configuration, 314–315
 patching inner nuclear membrane, 316
 studying single IP_3R channel activity
 in inner nuclear membrane, 303–304
 methods for, 298–300, 299f
 in outer nuclear membrane, 300–303, 301f, 302f
 reconstitution of endoplasmic reticulum IP_3R receptors into BLMs (protocol), 330–336
 materials, 330–331, 331f
 method, 331–334, 332f, 334f, 335f
 recipes, 336
 troubleshooting, 334–335
 regulation of, 156
 structure of, 156
 study using $^{45}Ca^{2+}$ -flux technique, 217, 222
 IP_3 . *See* Inositol 1,4,5-trisphosphate (IP_3)
 IP_3R . *See* Inositol 1,4,5-trisphosphate receptor (IP_3R)
 Institutional safety office, 571
 Intercellular Ca^{2+} wave (ICW)
 flash photolysis of caged IP_3 to trigger (protocol), 109–112
 materials, 109–110
 methods, 110–111
 ester loading of fluorescent Ca^{2+} indicator, 110
 quantification of ICW size, 111
 uncaging of IP_3 using flash photolysis, 110–111
 recipe, 112
 troubleshooting, 111–112
 functions of, 95–96
 induced by photolysis of electroporation-loaded caged IP_3 , 100f, 101
 mechanisms of propagation, 94–95, 94f
 methods to induce, 96
 in pathological conditions, 96
 quantification of ICW size, 111
 velocity of, 95
 Internal recording solution for Ca^{2+} currents (recipe), 269, 275
 Internal solution for calcium imaging (recipe), 149, 153
 Intracellular buffer (recipe), 191, 507
 Intracellular calcium (Ca^{2+})
 assessing Ca^{2+} -removal pathways in cardiac myocytes (protocol), 438–443
 discussion, 441–442
 materials, 438–439
 method, 439–441, 440f, 441f
 recipes, 442–443
 $^{45}Ca^{2+}$ fluxes in intact cells, technique of, 216–217
 $^{45}Ca^{2+}$ fluxes in permeabilized cells
 analysis of, 215–216, 216f
 technique, 213–215
 calcium signaling and, 277–278
 in excitation-contraction coupling, 411
 measurement of release, 211–229
 in intact cells using $^{45}Ca^{2+}$ (protocol), 225–229
 discussion, 227
 materials, 225–226
 method, 226–227
 recipes, 228
 troubleshooting, 227
 overview, 212–213
 in permeabilized cells using $^{45}Ca^{2+}$ (protocol), 219–224
 discussion, 221–222, 222f
 materials, 219
 methods, 219–221
 recipes, 222–223
 troubleshooting, 221
 models of signaling/dynamics, 553–569
 compartmentalized models, 555–557
 exploring oscillations in a point model of intracellular Ca^{2+} concentration (protocol), 562–565, 564f
 fire-diffuse-fire (FDF) models, 558–559, 559f
 ordinary differential equation (ODE) models, 554–555, 562–565
 spatially extended models, 557–559, 559f
 threshold models, 558
 time to blip-stochastic simulation of single channel opening (protocol), 566–569, 567f, 568f
 two-dimensional models, 558–559, 559f
 whole-cell models, 554–555
 monitoring yeast intracellular Ca^{2+} levels using *in vivo* bioluminescence assay (protocol), 456–459
 materials, 456–457
 method, 457–459, 458f
 troubleshooting, 459
 signaling, 211–212, 212f
 Intracellular Ca^{2+} waves, modeling calcium signaling and, 553–554, 557–559, 559f
 Intracellular medium (recipe), 515, 524
 Intracellular organelles, using targeted variants of aequorin to measure Ca^{2+} levels in (protocol), 185–191
 calibration, 187, 188f
 discussion, 191
 materials, 185–187, 186f
 measurement of Ca^{2+} in ER and its release, 190
 measurement of capacitative Ca^{2+} influx, 188–189
 measurement of cytosolic Ca^{2+} , 187–188
 measurement of mitochondrial Ca^{2+} uptake, 189, 189f
 measurement of mitochondrial Ca^{2+} uptake in permeabilized cells, 190
 recipes, 191
 reconstitution of functional photoprotein, 187
 transfection, 187
 troubleshooting, 190–191
 Intracellular solution with DM-Nitrophen and Fluo-4 (recipe), 119
 Inverted microscope, through-the-objective total internal reflection fluorescence (TIRF) microscopy on, 19
 Ion channels. *See also* Calcium (Ca^{2+}) channels
 investigating ion channel distribution using combination of spatially limited photolysis, Ca^{2+} imaging, and patch-clamp recording (protocol), 85–89
 studying activation of epithelial ion channels using global whole-field photolysis (protocol), 79–84
 Ion lasers, 12
 Ionomycin, 58, 62, 68, 190, 215, 216, 278
 IP_3 . *See* Inositol 1,4,5-trisphosphate (IP_3)
 IP_3R . *See* Inositol 1,4,5-trisphosphate receptor (IP_3R)
 IRBIT, 222
 Isoaequorins, 193
 Isosbestic point, 27
- ## J
- Jablonski energy transitions, 3
 Johnson, Frank H., 177
 JPW1114, 142f, 143, 146, 147f
 JPW-3028, 142
 Junctional sarcoplasmic reticulum, 411

K

- K-aspartate-based internal pipette solution (recipe), 436
- K-aspartate-based pipette solution (recipe), 418
- K⁺ current, 427
- Kilobeam array scanner, 116–120, 118f
- Krebs-Ringer Modified Buffer (KRB) (recipe), 191

L

- La³⁺, 287
- LACS (low-affinity calcium system), 446f, 447
- λ-protein phosphatase, 250
- Laminin, 401, 406
- Lampbrush chromosomes, 383
- Lasers
 - in confocal microscopy, 12
 - ion, 12
 - polarizing filters to attenuate laser power, 12
 - pulsed, 2
 - safety, 20, 572
 - stimulated emission depletion (STED) microscopy, 23
 - two-photon fluorescence and, 2–3
- Laser-scanning confocal microscopy (LSCM)
 - basic elements of, 8f
 - complications of, 9
 - dwell time, 9
 - image speed, 10–11, 10f
 - imaging Ca²⁺ dynamics in plants
 - high-resolution imaging of cytoplasmic Ca²⁺ dynamics in *Arabidopsis* roots (protocol), 474–478, 476f
 - imaging of mitochondrial and nuclear Ca²⁺ dynamics in *Arabidopsis* roots (protocol), 484–488, 485f
 - live cell imaging of cytoplasmic and nuclear Ca²⁺ dynamics in *Arabidopsis* roots (protocol), 479–483, 481f
 - lasers, 12
 - line scan, 11
 - measuring Ca²⁺ sparks in cardiac myocytes (protocol), 430–437
 - photomultiplier gain, 11–12
 - pinhole size, 9–10, 9f
 - principles of operation, 7–8, 8f
 - spatially limited photolysis, 77
 - spectral imaging, 12–13
- LDAO, voltage-dependent anion-selective channel protein 1 (VDAC1) purification using, 345
- LDAO solution for purification of VDAC1 from mitochondria (recipe), 348
- LED light sources, 5, 28
- Lens chromatic aberration, 7
- Letm1, 338, 339
- L-glutamate photorelease, combining Ca²⁺ imaging with (protocol), 150–153, 151f
 - configuration of imaging equipment, 151–152, 151f
 - materials, 150–151
 - photorelease, 152
 - recipes, 152–153
 - troubleshooting, 152
- Light sources, for spatially limited photolysis, 77
- Line scan, in confocal microscopy, 11
- Liposomes, assay of Ca²⁺ transport by VDAC1 reconstituted into (protocol), 355–358
 - Ca²⁺ efflux from VDAC1-containing proteoliposomes, 356–357
 - discussion, 357
 - liposome preparation, 356
 - materials, 355–356
 - method, 356–357
 - recipe, 357

- reconstitution of VDAC1 and loading with Ca²⁺, 356
- troubleshooting, 357
- Li-Rinzel approximation of the De Young-Keizer model, 562–565, 564f
- Loading medium for intact cells (recipe), 228
- Loading medium for permeabilized cells (recipe), 223
- Look-up table, 33
- Low-conductivity electroporation buffer (recipe), 107
- L-type Ca²⁺ current (*I*_{CaL}), 420–423
- L-type calcium channels (LTCCs), 411–412
- voltage-dependent Ca²⁺ channels (VDCCs), 555–556
- Lysis buffer for microsome preparation (recipe), 242
- Lysosensor Green DND-189, 521, 522
- Lysosensor Yellow/Blue DND-160, 521, 522
- Lysotracker Red DND-99, 521, 522

M

- Macspark, 435
- Mag-Fluo-4, in high-throughput functional assays of IP₃-evoked Ca²⁺ release protocol, 161, 163
- Mag-Fura-2, dissociation constant of, 141
- Malignant hyperthermia mutations, 323
- Manganese oxide, 526, 527
- Master mix (4×) (recipe), 242
- Master mix for Ca²⁺-uptake assay (recipe), 254
- Material safety data sheets (MSDSs), 571
- MathLab (software), 137
- MATLAB (software), 566–568
- Maxchelator (software), 82–83, 433, 537
- Media exchanges, 38–39
- Medical Pathological Waste (MPW), 573
- Membrane potential imaging
 - combining Ca²⁺ and V_M optical measurements, 141–143, 142f
 - combining with Ca²⁺ imaging in single neurons (protocol), 146–149
 - discussion, 149
 - materials, 146–147
 - method, 147–148, 147f
 - recipes, 149
 - troubleshooting, 148
 - overview, 139–140
- Mercury arc lamps, 28
- Metal-chelate buffer systems, 66–71
 - buffering ranges of, 67t
 - [Ca²⁺]_{free} of Ca²⁺-EDTA, 67t
 - [Ca²⁺]_{free} of Ca²⁺-EGTA, 68t
 - [Ca²⁺]_{free} of Ca²⁺-HEDTA, 68t
- MetaMorph (software), 404
- Methylnitro indoliny (MNI)-caged glutamate, 77
- Mg²⁺-free cytosol-like medium (recipe), 165
- Microinjection
 - of apoaequorin mRNA into zebrafish embryos, 209
 - of cyclic adenosine dinucleotide phosphate ribose (cADPR), 505–506
 - of holo-aequorin into dechorionated and intact zebrafish embryos (protocol), 197–205
 - of nicotinic acid adenine dinucleotide phosphate (NAADP), 523
- Microinjection pipettes, preparation of, 198, 199f
- Micro-Manager (software), 28
- Microscopy. *See* Fluorescence microscopy
- Microsomes
 - high-throughput measurement of Ca²⁺-dependent ATPase in (protocol), 234–244
 - preparation of microsomes to study Ca²⁺ channels (protocol), 326–329
 - harvesting rat cerebella, 327
 - materials, 326–327
 - method, 327–328
 - microsomal preparation, 328
 - preparation of Sf9 cells, 327–328
 - recipes, 329
 - troubleshooting, 328
- Microwave safety, 572
- Mid2p, 450
- Mid1 protein/gene, 446f, 448t, 449f, 450
- Minimal medium (recipe), 409
- Mirror, dichroic, 4, 5
- Mitochondria
 - calcium (Ca²⁺) transport (*see* Mitochondrial Ca²⁺ transport)
 - Ca²⁺ uptake, measurement of, 189, 189f
 - in permeabilized cells, 190
 - imaging of mitochondrial and nuclear Ca²⁺ dynamics in *Arabidopsis* roots (protocol), 484–488
 - isolation of, 344
 - voltage-dependent anion-selective channel protein 1 (VDAC1) purification from rat liver (protocol), 343–348
- Mitochondria Ca²⁺ uniporter (MCU), 337–339
- Mitochondrial calcium uptake 1 (MICU1), 339
- Mitochondrial Ca²⁺ transport, 337–366
 - assays of mitochondrial Ca²⁺ transport and Ca²⁺ efflux via MPTP (protocol), 359–361, 361f
 - discussion, 360, 361f
 - materials, 359–360
 - method, 360
 - recipe, 361
 - components of, 338–339, 338f
 - importance of mitochondrial calcium, 337–338
 - mitochondrial Na⁺/Ca²⁺ exchange assays (protocol), 362–366
 - discussion, 364
 - materials, 362–363
 - method, 363–364, 364f
 - recipes, 365–366
 - troubleshooting, 364
 - voltage-dependent anion-selective channel protein 1 (VDAC1), 337–341, 338f
 - assay of Ca²⁺ transport by VDAC1 reconstituted into liposomes, 355–358
 - Ca²⁺ efflux from VDAC1-containing proteoliposomes, 356–357
 - discussion, 357
 - liposome preparation, 356
 - materials, 355–356
 - method, 356–357
 - recipe, 357
 - reconstitution of VDAC1 and loading with Ca²⁺, 356
 - troubleshooting, 357
- channel conductance and permeability, 340
- conformational states, 340
- permeability to Ca²⁺, 340–341
- purification from rat liver mitochondria (protocol), 343–348
 - discussion, 346
 - isolation of mitochondria, 344
 - materials, 343–344
 - method, 344–346
 - recipes, 346–348
 - troubleshooting, 346
 - VDAC1 purification using carboxymethyl (CM)-cellulose, 345–346
 - VDAC1 purification using LDAO, 345
 - VDAC1 purification using reactive red-agarose (RRA), 345
 - VDAC1 purification using Triton X-100, 344–345

- reconstitution of purified VDAC1 into lipid bilayer and recording of channel conductance (protocol), 349–354
assembly of bilayer system, 350–351
discussion, 353
formation of planar bilayer membrane, 351
materials, 349–350
method, 350–353, 351f
recipes, 353
reconstitution of VDAC1 into planar lipid bilayer, 351–352
recording currents through VDAC1, 352–353, 352f
troubleshooting, 353
- Mitochondrial inhibitors, 215
- Mitochondrial ratiometric pericam (RP-mt), 363, 364t
- Mitochondrial ryanodine receptor (mRyR), 339
- Mitochondria NCLX (Na⁺/Ca²⁺ exchanger), 337–339, 364
- Mitochondria permeability transition pore (MPTP)
assays of mitochondrial Ca²⁺ transport and Ca²⁺ efflux via MPTP (protocol), 359–361, 361f
discussion, 360, 361f
materials, 359–360
method, 360
recipe, 361
described, 337–339, 343
- MitoTracker Red, 57
- MLIV (mucopolipidosis type IV), 447
- mMESSAGE mMACHINE SP6 kit, 208
- Mnemioptin, 177
- MNI (4-methoxy-7-nitroindolyl)-caged L-glutamate, 143–144, 150–152, 151f
- Modified Barth's saline (MBS) (1×, pH 7.8) (recipe), 393
- MonoQ columns, 527
- Mpk1 MAP kinase cascade, 449
- MPTP. *See* Mitochondria permeability transition pore
- MPW (Medical Pathological Waste), 573
- mRNA injection into zebrafish embryos, 209
- mRyR (mitochondrial ryanodine receptor), 339
- MSDSs (material safety data sheets), 571
- MS medium with sucrose (liquid) (recipe), 483
- MS medium with sucrose (solid) (recipe), 483
- Mucopolipidosis type IV (MLIV), 447
- Multichannel imaging of cellular signaling (protocol), 126–129
discussion, 129
materials, 126–127
method, 127–128, 128f
recipe, 129
troubleshooting, 128–129
- Myocytes. *See* Cardiomyocytes
- N**
- NA (numerical aperture), objective lens, 30–31, 31t
- NAADP. *See* Nicotinic acid adenine dinucleotide phosphate
- NAADP-AM (nicotinic acid adenine dinucleotide phosphate-acetoxymethyl ester), 517–520, 518f
- Na⁺-containing solution for measuring Ca²⁺ fluxes across the mitochondrial inner membrane (recipe), 365
- NAD (nicotinamide adenine dinucleotide), 490, 491f, 493, 495, 496, 499, 530–531
- NADH (nicotinamide adenine dinucleotide hydrogen), 493
- NADP (nicotinamide adenine dinucleotide phosphate), 509, 510f
- NAD reaction buffer (recipe), 531
- Na⁺-free solution for measuring Ca²⁺ fluxes across the mitochondrial inner membrane (recipe), 365
- NaN₃, 215
- National Institute of Environmental Health and Human Services, 575
- NBQX, 144
- NCLX. *See* Sodium/calcium (Na⁺/Ca²⁺) exchanger
- NCX. *See* Sodium/calcium (Na⁺/Ca²⁺) exchanger
- n*-decane, 350
- NEP-InsP₃, 79, 81f, 82
- Nernst equation, 368
- Neurons
calcium transients in developing *Xenopus* spinal neurons (protocol), 399–409
discussion, 407–408
materials, 399–400
method, 400–407
data analysis, 404
general imaging methodology, 401–402, 402f
imaging filopodial transients in vivo, 403–404, 403f
imaging growth cone transients in vivo, 402–403, 403f
imposition of Ca²⁺ spikes, 405, 405f
manipulation of Ca²⁺ dynamics in growth cones, 406
manipulation of filopodial Ca²⁺ dynamics, 406–407
perturbation experiments, 404
preparation of dissociated cell cultures, 400–401
preparation of spinal cords, 401
recipes, 408–409
troubleshooting, 407
combining Ca²⁺ and membrane potential imaging in single neurons (protocol), 146–149, 147f
dorsal root ganglion (DRG) neurons
dissecting, 265, 266f
L-type Ca²⁺ current in, 260
N-type Ca²⁺ current in, 260
whole-cell patch-clamp recordings of Ca²⁺ currents from isolated neonatal dorsal root ganglion (DRG) neurons (protocol), 264–270
- NiCl₂, 438–439, 440f, 441
- Nicolson-Eisenman equation, 368
- Nicotinamide, 493, 495, 498
- Nicotinamide adenine dinucleotide (NAD), 490, 491f, 493, 495, 496, 499, 530–531
- Nicotinamide adenine dinucleotide hydrogen (NADH), 493
- Nicotinamide adenine dinucleotide phosphate (NADP), 509, 510f
- Nicotinic acid, 530
- Nicotinic acid adenine dinucleotide phosphate (NAADP), 212–213, 509–531
overview of, 509–510
protocols, 512–531
caged NAADP, 525–528
discussion, 528
materials, 525–526
method, 526–527
troubleshooting, 527–528
measurement of luminal pH of acidic stores as a readout for NAADP action, 521–524
discussion, 523
materials, 521–522
method, 522–523
recipes, 523–524
preparation and use of sea urchin egg homogenates for studying NAADP-mediated Ca²⁺ release, 512–516
materials, 512–513
- method, 513–514, 514f
recipes, 515
troubleshooting, 514
synthesis of NAADP-AM as a membrane-permeant NAADP analog, 517–520
discussion, 519–520
materials, 517–518
method, 518–519, 518f
troubleshooting, 519
synthesis of [³²P]NAADP for radioreceptor binding assay, 529–531, 530f
discussion, 530–531
materials, 529
method, 530, 530f
recipe, 531
structure of, 510f
- Nicotinic acid adenine dinucleotide phosphate-acetoxymethyl ester (NAADP-AM), 517–520, 518f
- Nifedipine, 267, 439, 447
- Nifedipine stock solution (recipe), 270
- Nisoldipine, 421
- Nitrilo triacetic acid (NTA), affinity for Ca²⁺, 537t
- 2-nitroacetophenone hydrazone, 526
- Nitrobenzofuran-EGTA, 77
- NMDA receptors, 142
- Nominally Ca²⁺-free artificial seawater (recipe), 515
- Norepinephrine, 188
- NP-EGTA, 79, 81f, 85–86, 86f, 88
- NP-EGTA-AM, 90–91
- NTA (nitrilo triacetic acid), affinity for Ca²⁺, 537t
- Nuclear Ca²⁺ dynamics
imaging of mitochondrial and nuclear Ca²⁺ dynamics in *Arabidopsis* roots (protocol), 484–488
live cell imaging of cytoplasmic and nuclear Ca²⁺ dynamics in *Arabidopsis* roots (protocol), 479–483
- Nuclear microinjection to assess impact of heterologously expressed proteins on Ca²⁺ signals in *Xenopus* oocytes (protocol), 388–394
- Nuclear patch-clamping
isolating nuclei from cultured cells (protocol), 308–312, 309t, 311f
discussion, 311
materials, 308–309, 309t
method, 309–311, 311f
recipe, 312
nuclear patch-clamp protocol, 313–318
cytoplasmic-side-out configuration, 315–316
luminal-side-out configuration, 315
materials, 313–314
on-nucleus configuration, 314–315
patching inner nuclear membrane, 316
studying single InsP₃R channel activity in inner nuclear membrane, 303–304, 304f
methods for, 298–300, 299f
in outer nuclear membrane, 300–303, 301f, 302f
studying single RyR channels in outer nuclear membrane, 304
- Nuclease-free H₂O (recipe), 210
- Nucleotide pyrophosphatase, 495
- Numerical aperture (NA), objective lens, 30–31, 31t
- Nystatin, 280, 418
- O**
- Obeline, 177
- Objective lens
imaging Ca²⁺ indicators and, 30–31, 31t
in through-the-objective total internal reflection fluorescence (TIRF) microscopy, 18–19, 19f

- Occupational Safety and Health Administration (OSHA), 571
- Offset control, 32f, 33
- OGB-5N (Oregon Green BAPTA-5N), 144, 144f, 536, 539, 539t, 546–550, 547f
- ω -agatoxin IVA, 260
- ω -conotoxin GVIA stock solution (recipe), 270
- ω -conotoxin GVIA, 260, 267
- One-photon fluorescence, 2–3, 2f
- Oocyte, of *Xenopus*
mammalian cell compared, 384t
as single-cell model for studying Ca^{2+} signaling, 381–398
- Oocyte dissociation solution (recipe), 393
- Optical parametric oscillator (OPO), 15
- ORAI channels
overview, 277–281, 280f
protocols
measuring endogenous I_{CRAC} and ORAI currents with the patch-clamp technique, 283–290, 287f
minimum requirements to use calcium imaging to analyze I_{CRAC} , 291–295, 293f
- Ordinary differential equation (ODE) models, of intracellular calcium signaling, 554–555, 562–565
- Oregon Green, 140, 144
- Oregon Green BAPTA-5N, 144, 144f, 536, 539, 539t, 546–550, 547f
- OSHA (Occupational Safety and Health Administration), 571
- ## P
- PALM/STORM super-resolution imaging, 21–23
single particle tracking (sptPALM), 22
- Pannexins (Panxs), 94, 95
- Parallel fiber-Purkinje neuron (PF-PN) synapses, 143
- Patch-clamp technique, 157
in cardiac myocytes, 416–417
described, 257–259, 258f
electrophysiology of intracellular Ca^{2+} channels, 297–319
isolating nuclei from cultured cells (protocol), 308–312, 309t, 311f
discussion, 311
materials, 308–309, 309t
method, 309–311, 311f
recipe, 312
nuclear patch-clamp protocol, 313–318
cytoplasmic-side-out configuration, 315–316
luminal-side-out configuration, 315
materials, 313–314
on-nucleus configuration, 314–315
patching inner nuclear membrane, 316
overview, 297
studying single InsP_3R channel activity in inner nuclear membrane, 303–304
methods for, 298–300, 299f
in outer nuclear membrane, 300–303, 301f, 302f
studying single RyR channels in outer nuclear membrane, 304, 304f
- error sources, 268–269
- investigating ion channel distribution using combination of spatially limited photolysis, Ca^{2+} imaging, and patch-clamp recording (protocol), 85–89
- measurement of I_{CRAC} and ORAI channel activity
overview, 277–281, 280f
protocols
measuring endogenous I_{CRAC} and ORAI currents with the patch-clamp technique, 283–290, 287f
minimum requirements to use calcium imaging to analyze I_{CRAC} , 291–295, 293f
- perforated patch, 280, 280f, 418
- recording voltage-sensitive Ca^{2+} channels, 257–276
current isolation methods, 268
overview, 257–262
whole-cell recordings of Ca^{2+} currents
from isolated neonatal dorsal root ganglion (DRG) neurons (protocol), 264–270
discussion, 268–269
materials, 264–265
method, 265–267
recipes, 269–270
troubleshooting, 267–268
- whole-cell recordings of recombinant channels heterologously expressed in HEK-293 cells (protocol), 271–275, 273f
discussion, 274
materials, 271–272
method, 272–274, 273f
recipes, 275
troubleshooting, 274
- Patch pipette solution for InsP_3R (recipe), 318
- Patch pipette solution for RyR (recipe), 318
- PCA (perchloric acid) extraction, 494, 495, 496
- Pearson's correlation analysis, 57
- Perchloric acid (PCA) extraction, 494, 495, 496
- Pericams
binding parameters of, 52t
mitochondrial ratiometric pericam (RP-mt), 363, 364t
- Permeability transition pore (PTP), 337–339, 338f. *See also* Mitochondria permeability transition pore
- Permeabilization solution for measuring Ca^{2+} fluxes across the mitochondrial inner membrane (recipe), 366
- Permeabilized-cell internal solution (350 μM EGTA) (recipe), 437
- PF-PN (parallel fiber-Purkinje neuron) synapses, 143
- pH
measurement of luminal pH of acidic stores as a readout for NAADP action (protocol), 521–524
discussion, 523
materials, 521–522
method, 522–523
recipes, 523–524
pH buffers, pK_a and temperature dependent of common, 542t
- PHERAstar plate-reader, 168–170
- Pheromones, yeast and, 447, 450
- Phialidin, 177
- Phosphatidylcholine, 331–332
- Phosphatidylethanolamine, 331–332
- Phosphatidylinositol bisphosphate (PIP_2), 94, 322, 450, 451f
- Phosphatidylserine, 331–332
- Phosphocreatine, 505, 513, 522
- Phospholamban, 241, 245, 250–253, 250f
- Phospholipase C
 IP_3 regeneration by, 94, 94f, 95
isoforms of, 94
- Phosphorylation mix ($4\times$) (recipe), 243
- Photoactivated localization microscopy (PALM), 21–23
- Photobleaching
 Ca^{2+} indicators choice and, 29–30
with confocal microscopy, 11
with two-photon microscopy, 15
with wide-field fluorescence microscopy, 6
- Photolysis
combining Ca^{2+} imaging with L-glutamate photorelease (protocol), 143–144, 144f, 150–153, 151f
in determining the Ca^{2+} -binding properties of DM-nitrophen and proteins (protocol), 546–551, 547f
flash photolysis of caged IP_3 to trigger intercellular Ca^{2+} waves (protocol), 109–112
in measuring rate constants of Ca^{2+} indicators (protocol), 541–545
two-photon photolysis combined with kilobeam array scanner to probe calcium signaling in cardiomyocytes (protocol), 116–120
two-photon photolysis of MNI-glutamate, 144
- Photomultiplier gain, 11–12
- Photomultiplier tube (PMT)
avalanche, 11
in confocal microscopy, 11–12
InGaAsP, 11
in two-photon microscopy, 14, 14f
- Photoprotein, 177. *See also* Aequorin
- Photorelease, combining Ca^{2+} imaging with L-glutamate (protocol), 143–144, 144f, 150–153, 151f
- Phototoxicity
 Ca^{2+} indicators choice and, 29–30
with confocal microscopy, 11
with flash photolysis of caged IP_3 , 112
with wide-field fluorescence microscopy, 6
- Pinhole size, in confocal microscopy, 9–10, 9f
- PIP_2 . *See* Phosphatidylinositol bisphosphate
- Pipettes, preparation of microinjection, 198, 199f
- Pipette solution for Cl^- current recordings (recipe), 83
- Pipette solution for K^+ current recordings (recipe), 84, 88
- Pixel binning, 31
- Pixel registration, problem with wide-field fluorescence microscopy, 6–7
- Planar lipid bilayer (PLB). *See also* Black lipid membrane (BLM)
reconstitution of purified VDAC1 into lipid bilayer and recording of channel conductance (protocol), 349–354, 351f, 352f
- Plants, Ca^{2+} imaging in, 465–488
 Ca^{2+} dynamics, 467–468
high-resolution imaging of cytoplasmic Ca^{2+} dynamics in *Arabidopsis* roots (protocol), 474–478
discussion, 477
materials, 474–475
method, 475–477, 476f
data analysis, 477
imaging, 475–477
plant growth, 475
sample preparation, 475, 476f
troubleshooting, 477
- imaging of mitochondrial and nuclear Ca^{2+} dynamics in *Arabidopsis* roots (protocol), 484–488
discussion, 487
materials, 484–485
method, 485–486
CLSM analysis, 486
imaging analyses, 486
sample preparation, 486
seed sterilization and growth of plantlets, 485–486
recipes, 487
troubleshooting, 487
- live cell imaging of cytoplasmic and nuclear Ca^{2+} dynamics in *Arabidopsis* roots (protocol), 479–483

- discussion, 482
materials, 479–480
method, 480–482, 481f
Ca²⁺ imaging, 480
data analyses, 482
plant growth, 480
sample preparation, 480, 481f
recipes, 483
troubleshooting, 482
- live cell imaging of cytoplasmic Ca²⁺ dynamics in *Arabidopsis* guard cells (protocol), 469–473
discussion, 473
materials, 469–470
method, 470–472, 471f
data analysis, 472
imaging, 471–472
plant growth, 470
sample preparation, 470–471, 471f
recipe, 473
troubleshooting, 472–473
- overview, 465–486
Yellow Cameleon Ca²⁺ indicators, 466–467, 466f
- Plasma membrane Ca²⁺ ATPase (PMCA), 231, 277, 291, 293, 438, 446
- PLB. *See* Planar lipid bilayer
- Plc1, 450
- Pluronic F-127, 42–43, 110, 161, 519
- PMCA. *See* Plasma membrane Ca²⁺ ATPase
- Pmcl protein/gene, 446f, 447, 448t, 449f, 450, 451
- PMR2 gene, 450
- Pmr1 protein/gene, 446, 446f, 448t, 449f, 450
- PMT. *See* Photomultiplier tube
- Polycystins, 222
- Poly-D-lysine, 401
- Poly-L-lysine, 37, 160, 505
- Poly-L-lysine stock solution (recipe), 270, 275
- Porin 1. *See* Voltage-dependent anion-selective channel protein 1 (VDAC1)
- Potassium-based Ringer's solution for *Xenopus* neurons (recipe), 409
- Prism-based total internal reflection fluorescence (TIRF) microscopy, 18
- Probenecid, 45
- Progesterone, 391
- Programmed cell death, 451–452
- Promoters
GCI, 469
ubiquitin, 474
- Protein concentration, determination of, 249–250
- Protein kinase C, 126–129, 128f, 450
- Proteins, Ca²⁺-binding kinetics of, 533–551
challenges in studying, 533–534
collecting data to determine the Ca²⁺-binding properties of DM-nitrophen and proteins (protocol), 546–551
discussion, 549–550
materials, 546
method, 547–549, 547f
recipe, 550
troubleshooting, 549
- measuring rate constants of Ca²⁺ indicators (protocol), 541–545
discussion, 545
materials, 541–542, 542t, 543f
method, 543–544
recipe, 545
troubleshooting, 544–545
- measuring steady-state properties of Ca²⁺ indicators with a set of calibrated [Ca²⁺] solutions (protocol), 536–540
discussion, 539–540, 539t
materials, 536
method, 537–538, 537t, 538t
recipe, 540
- troubleshooting, 538
overview of method for measuring, 534–535, 534f
- Proteoliposomes, Ca²⁺ efflux from VDAC1-containing, 356–357
- Protonophores, 450
- PTP. *See* Permeability transition pore
- Pulsed laser, 2
- PVC use in Ca²⁺-sensitive electrodes, 369–370, 373, 377
- P2Y purinoceptors, 519
- ## Q
- QuickPALM (software), 23
- Quin-2, 367
- ## R
- Radiation
safety procedures, 574
waste disposal, 573, 574
- Radioreceptor binding assay, synthesis of [³²P]NAADP for (protocol), 529–531, 530f
discussion, 530–531
materials, 529
method, 530, 530f
recipe, 531
- RapidSTORM project, 23
- Rapid western blotting protocol *Xenopus* oocyte, 395–398
- Ratiometric indicators, 26–27
- Rayleigh criterion, 6
- Reactive oxygen species (ROS), 29, 30
- Reactive red-agarose (RRA), 345
- Reagent 1 (recipe), 243
- Reagent 2 (recipe), 243
- Recipes
ACSF for calcium imaging, 149, 152
aequorin buffer, 204
agarose solution for building agar salt bridges, 353
artificial seawater (ASW), 515, 523
ATP stock solution for ER sensor calibrations, 63
bath solution for InsP₃R, 317
bath solution for recording Cl⁻ currents, 83, 88
bath solution for recording K⁺ currents, 83, 88
bath solution for RyR, 318
buffer A for assaying Ca²⁺ transport by VDAC1 reconstituted into liposomes, 357
buffer A for isolation of mitochondria from rat liver, 346
buffer B for isolation of mitochondria from rat liver, 346
buffer B for microinjection, 507
buffer C for purification of VDAC1 from mitochondria, 347
buffer D for purification of VDAC1 from mitochondria, 347
buffer E for purification of VDAC1 from mitochondria, 347
buffer F for purification of VDAC1 from mitochondria, 347
buffer for assaying Ca²⁺ transport by VDAC1 reconstituted into PLBs, 353
buffer G for purification of VDAC1 from mitochondria, 347
buffer H for purification of VDAC1 from mitochondria, 347
buffer I for purification of VDAC1 from mitochondria, 348
Ca²⁺-containing modified Krebs solution, 228
CaEGTA (100 mM), 436
Ca-EGTA solution, 336
Ca²⁺-free cytosol-like medium, 165
- Ca²⁺-free perfusion solution, 124
calcium blocker cocktail, 408
calcium-chelator solution for metal-chelate buffer system, 72
calcium-free Ringer's solution for *Xenopus* neurons, 408
Ca²⁺ loading solution for measuring Ca²⁺ fluxes across the mitochondrial inner membrane, 365
Ca²⁺ mix, 242
Ca²⁺ uptake solution for assaying Ca²⁺ transport in mitochondria, 361
chelator solution for metal-chelate buffer system, 72
Chelex-treated ultrapure H₂O, 63, 72
cis recording solution, 336
Cs-aspartate-based internal pipette solution, 423, 428
CsCl-based internal pipette solution, 442
Cs-Tyrode-based external solution, 423, 429, 443
culture medium for cells expressing recombinant Ca²⁺ channels, 275
culture medium for DRG neurons, 269
Danieau's solution, 204, 210
dialysis buffer for purification of VDAC1 from mitochondria, 348
dissociating medium for DRG neurons, 269
divalent-cation-free minimal medium, 408
efflux medium for intact cells, 228
efflux medium for permeabilized cells, 222
EGTA (0.5 M), 63, 72
EGTA Ca²⁺-free artificial seawater, 515
EGTA mix, 242
EGTA solution (100 mM), 436
external recording solution for Ca²⁺ currents, 269
external recording solution for recombinant Ca²⁺ channels, 275
extracellular buffer, 501
extracellular solution with 1 mM Ca²⁺, 294
extracellular solution with no free Ca²⁺, 295
Fluo-3-AM solution, 409
Fluo-4-AM solution, 409
growth medium (*Arabidopsis*), 487
Hanks balanced salt solution-HEPES, 107, 112
HEPES-buffered Hanks balanced salt solution (HHBSS) with Ca²⁺, 64, 72
HEPES-buffered Hanks balanced salt solution without Ca²⁺ (Ca²⁺-free HHBSS), 64
HEPES-buffered Hanks balanced salt solution without Ca²⁺ and Mg²⁺ (Ca²⁺/Mg²⁺-free HHBSS), 73
HEPES-buffered saline (HBS), 165
high-salt buffer (HSB) for microsomal preparation, 329
homogenization buffer for Ca²⁺-uptake assay, 253
homogenization buffer (HB) for cerebellar microsomal preparation, 329
homogenization buffer for sea urchin homogenates, 515
homogenization buffer (SF9HB) for Sf9 microsomal preparation, 329
HPLC buffer A, 501
HPLC buffer B, 501
I_{CRAC} Ca²⁺-containing bath solution, 289
I_{CRAC} Ca²⁺-free bath solution, 289
I_{CRAC} divalent-free (DVF) bath solution, 289
I_{CRAC} pipette solution for active depletion, 289
I_{CRAC} pipette solution for passive depletion, 289
imaging medium (*Arabidopsis*), 487
imaging solution, 92
incubation buffer, 473
internal recording solution for Ca²⁺ currents, 269, 275
internal solution for calcium imaging, 149, 153

Recipes (Continued)

- intracellular buffer, 191, 507
intracellular medium, 515, 524
intracellular solution with DM-Nitrophen and Fluo-4, 119
K-aspartate-based internal pipette solution, 436
K-aspartate-based pipette solution, 418
Krebs-Ringer Modified Buffer (KRB), 191
LDAO solution for purification of VDACL from mitochondria, 348
loading medium for intact cells, 228
loading medium for permeabilized cells, 223
low-conductivity electroporation buffer, 107
lysis buffer for microsome preparation, 242
master mix (4×), 242
master mix for Ca²⁺-uptake assay, 254
Mg²⁺-free cytosol-like medium, 165
minimal medium, 409
modified Barth's saline (MBS) (1×, pH 7.8), 393
MS medium with sucrose (liquid), 483
MS medium with sucrose (solid), 483
Na⁺-containing solution for measuring Ca²⁺ fluxes across the mitochondrial inner membrane, 365
NAD reaction buffer, 531
Na⁺-free solution for measuring Ca²⁺ fluxes across the mitochondrial inner membrane, 365
nifedipine stock solution, 270
nominally Ca²⁺-free artificial seawater, 515
nuclease-free H₂O, 210
ω-conotoxin GVIA stock solution, 270
oocyte dissociation solution, 393
patch pipette solution for InsP₃R, 318
patch pipette solution for RyR, 318
permeabilization solution for measuring Ca²⁺ fluxes across the mitochondrial inner membrane, 366
permeabilized-cell internal solution (350 μM EGTA), 437
phosphorylation mix (4×), 243
pipette solution for Cl⁻ current recordings, 83
pipette solution for K⁺ current recordings, 84, 88
poly-L-lysine stock solution, 270, 275
potassium-based Ringer's solution for *Xenopus* neurons, 409
reagent 1, 243
reagent 2, 243
recording solution for measuring Ca²⁺ fluxes across the mitochondrial inner membrane, 366
Ringer's solution for measuring Ca²⁺ fluxes across the mitochondrial inner membrane, 366
Ringer's solution for *Xenopus* neurons, 409
sample buffer for phosphorylation assay, 243
S⁺ and S⁻ media, 223
solution 1M, 243
standard reaction buffer for Ca²⁺ indicators, 540, 545, 550
stop solution for phosphorylation assay, 244
storage buffer for microsomal preparation, 329
sucrose buffer, 312
TBST for western blotting, 398
tetrodotoxin (TTX) stock solution, 270
trans recording solution, 336
Tris-acetate-EDTA (TAE) buffer (50×), 210
Triton X-100 solution for purification of VDACL from mitochondria, 348
trypsin solution for HEK-293 cells, 275
Tyrode's external solution, 437, 443
Tyrode solution for bath perfusion, 418, 424, 429
Tyrode solution for probing Ca²⁺ signaling, 120, 124, 129, 133, 138
vegetable oil (EGTA-washed), 204
wash buffer for Ca²⁺-uptake assay, 254
- Recording solution for measuring Ca²⁺ fluxes across the mitochondrial inner membrane (recipe), 366
- Red light, 2–3
Resazurin, 493, 495
Research Genetics, 445
Reverse-phase HPLC, measuring CD38 activity by (protocol), 498–502
discussion, 500–501
materials, 498–499
method, 499–500
recipes, 501
troubleshooting, 500
- Rhod-FF, 539
Ringer's solution for measuring Ca²⁺ fluxes across the mitochondrial inner membrane (recipe), 366
Ringer's solution for *Xenopus* neurons (recipe), 409
RNase Block, 208
RNaseZap, 208
RNA transcription, 208
Rolling ball subtraction method, 35, 36f
- Roots, *Arabidopsis*
high-resolution imaging of cytoplasmic Ca²⁺ dynamics in (protocol), 474–478
imaging of mitochondrial and nuclear Ca²⁺ dynamics in (protocol), 484–488
live cell imaging of cytoplasmic and nuclear Ca²⁺ dynamics in (protocol), 479–483
- ROS (reactive oxygen species), 29, 30
RP-mt (mitochondrial ratiometric pericam), 363, 364t
- RRA (reactive red-agarose), 345
Ru360, 339
Ruthenium red, 334, 339, 356
Ryanodine receptor (RyR), 94, 212, 277
caffeine effect on, 426–428, 427f, 433, 433f
in cardiac myocytes, 411–413, 426–428, 430–436
cyclic adenosine dinucleotide phosphate ribose (cADPR) and, 489–490, 491f
measuring Ca²⁺ sparks in cardiac myocytes (protocol), 430–437
analysis of Ca²⁺-spark recordings, 435
discussion, 434–436
evaluating ryanodine receptor function in intact cells, 431–432
evaluating ryanodine receptor function in permeabilized cells, 433–434, 433f
intact *versus* permeabilized cells, 434–435
materials, 430–431
method, 431–434, 433f
recipes, 436–437
SR Ca²⁺ leak, 435
mitochondrial (mRyR), 339
in models of intracellular calcium signaling, 554–556, 558
- Ryanodine receptor (RyR) channels
black lipid membrane (BLM) experiments to study InsP₃R and RyanR, 322–323
channels in endoplasmic reticulum, 321–323
patch-clamp study of single channels in outer nuclear membrane, 304, 304f, 313
- RyanR. *See* Ryanodine receptor; Ryanodine receptor (RyR) channels
- RyR. *See* Ryanodine receptor; Ryanodine receptor (RyR) channels
- ## S
- Saccharomyces cerevisiae*, 445–452, 446f, 448t, 449f.
See also Yeast
- Saccharomyces* Genome Database (SGD), 445
- Safety, 571–576
biological safety procedures, 574–575
general cautions, 571–573
institutional safety office, 571
- laser, 20
material safety data sheets (MSDSs), 571
properties of common hazardous chemicals, 575–576
radioactive safety procedures, 574
waste disposal, 573
- Sample buffer for phosphorylation assay (recipe), 243
- Sample handling
mounting, 37–38, 38f
perfusion, 38, 38f
solution exchange, 38–39
- S⁺ and S⁻ media (recipe), 223
- Saponin, 58, 63, 161, 213, 219, 433, 504, 506
- Sarco-endoplasmic Ca²⁺ ATPases (SERCAs), 180, 211, 214–215, 222, 277, 291, 446
measuring activity of, 231–255
measuring calcium (Ca²⁺) pump activity high-throughput measurement of Ca²⁺-dependent ATPase in COS microsomes (protocol), 234–244
measuring Ca²⁺-dependent Ca²⁺-uptake activity in the mouse heart (protocol), 245–254
mechanisms and features of action, 231–232
strategies for measuring transporter activity, 232–233
models of, 556
in myocytes, 411, 438, 441–442
- Sarcoplipin, 241, 245, 252
- Sarcoplasmic reticulum
calcium release from, 411, 413, 418
characterizing trigger in cardiac myocytes (protocol), 420–424, 422f
measuring fractional in cardiac myocytes (protocol), 425–429
junctional, 411
measuring Ca²⁺ content, 425–428, 427f
- Sea urchins
egg collection, 513
measurement of luminal pH of acidic stores as a readout for NAADP action (protocol), 521–524
discussion, 523
materials, 521–522
method, 522–523
recipes, 523–524
preparation and use of sea urchin egg homogenates for studying NAADP-mediated Ca²⁺ release (protocol), 512–516
materials, 512–513
method, 513–514, 514f
recipes, 515
troubleshooting, 514
- Second-harmonic imaging, 16
- Second messenger, criteria for, 489
- Second-messenger-operated channels (SMOCs), 278
- Secretory pathway Ca²⁺ ATPase (SPCA), 231, 446
- Selective illumination, 28–29
- SERCA. *See* Sarco-endoplasmic Ca²⁺ ATPases
- Sf9 cells, preparation of, 327–328
- Shimomura, Osamu, 177–178
- Signaling. *See* Calcium (Ca²⁺) signaling
- Simon, Willy, 369
- Single-wavelength Ca²⁺ indicators, 27–28
- Slr2, 450
- SMOCs (second-messenger-operated channels), 278
- Snf3, 451, 451f
- SOCCs (store-operated calcium channels), 278, 278f
- SOCE (store-operated calcium entry), 278, 278f, 279
- Sodium/calcium (Na⁺/Ca²⁺) exchanger, 291. *See also* Mitochondria NCLX (Na⁺/Ca²⁺ exchanger)

- in cardiac myocytes, 412, 413, 421–423, 438–442
 current through, 425–428, 427f, 438–441, 440f, 441f
 mitochondrial, 337–339, 364
 Sodium (Na^+) current, 423
 Sodium (Na^+) fluxes across mitochondrial inner membrane, measuring, 363–364, 364f
 Sodium/lithium (Na^+/Li^+) transporter, 448t, 450
 Sodium pentobarbital, 246
 Softmax Pro, 163
 SOICR (store-overload-induced Ca^{2+} release), 323
 Solution 1M (recipe), 243
 Solvents, safe handling of, 572, 575
 Soybean asolectin, 350, 356
 Spatially limited photolysis
 analyzing Ca^{2+} dynamics in intact epithelial cells using spatially limited flash photolysis (protocol), 90–92, 91f
 investigating ion channel distribution using combination of spatially limited photolysis, Ca^{2+} imaging, and patch-clamp recording (protocol), 85–89
 light sources for, 77
 SPCA (secretory pathway Ca^{2+} ATPase), 231, 446
 SPC-Image (software), 137
 Spectral imaging, in confocal microscopy, 12–13
 Spinning disc confocal microscopy, 13
 Standard reaction buffer for Ca^{2+} indicators (recipe), 540, 545, 550
 STED. *See* Stimulated emission depletion (STED) microscopy
 STIM proteins, 277, 278f, 279, 287, 287f, 288
 Stimulated emission depletion (STED) microscopy, 3, 23, 23f
 Stochastic optical reconstruction microscopy (STORM), 21–23
 Stokes shifts, 3
 Stop solution for phosphorylation assay (recipe), 244
 Storage buffer for microsomal preparation (recipe), 329
 Storage of image, 33–34
 Store-operated calcium channels (SOCCs), 278, 278f
 Store-operated calcium entry (SOCE), 278, 278f, 279
 Store-overload-induced Ca^{2+} release (SOICR), 323
 Stress response in yeast, 448, 450
 Styrl voltage-sensitive dyes, 140, 141
 Subtraction of background signal, 34, 35, 35f, 36f, 39
 Sucrose buffer (recipe), 312
 Sugar-induced calcium signaling pathway, 450–451, 451f
 Sulfinpyrazone, 45
 Super-resolution microscopy, 21–23
 PALM/STORM super-resolution imaging, 21–23
 STED super-resolution imaging, 23, 23f
 Swe1, 450
 SYLGARD 184, 56, 57f
 SymPhoTime (software), 137
- T**
- tBHQ, 292
 TBST for western blotting (recipe), 398
 TCA (tricarboxylic acid) cycle, 338, 338f
 T cells (T lymphocytes), 277, 278f, 281, 291–294, 503–506
 Tenascin, 401
 Tetrahydrofuran use in Ca^{2+} -sensitive electrodes, 373–374, 377
 Tetrodotoxin (TTX), 144, 423, 439
 Tetrodotoxin (TTX) stock solution (recipe), 270
 Thapsigargin, 59, 161, 211, 215, 216f, 241, 278, 286, 292, 439
 Through-the-objective total internal reflection fluorescence (TIRF) microscopy, 18–19, 19f
 TIRF. *See* Total internal reflection fluorescence (TIRF) microscopy
 Ti:Sapphire lasers, 15
 Tissue homogenization
 of mouse heart, 247–248
 for nuclei isolation, 310, 311
 of rat liver, 344
 of sea urchin eggs, 513
 TN-XL, 134–138, 136f
 Topological paradox, 490
 Total internal reflection fluorescence (TIRF) microscopy
 blocking filters for, 4–5
 overview, 17–21
 imaging, 20–21
 prism-based, 18
 safety issues, 20
 sample preparation, 21
 through-the-objective, 18–19, 19f
 selective illumination, 28–29
 Toxic compounds, 576
 Trans-Blot Turbo system, 398
 Transient receptor potential (TRP) channels, 277, 278
 Trans-illumination imaging
 in total internal reflection fluorescence (TIRF) microscopy, 18
 in two-photon microscopy, 16–17, 17f
 Transport proteins, mitochondrial, 337–341.
 See also specific transport proteins
 Trans recording solution (recipe), 336
 Tricarboxylic acid (TCA) cycle, 338, 338f
 Tris-acetate-EDTA (TAE) buffer (50 \times) (recipe), 210
 Triton X-100, 202, 203
 voltage-dependent anion-selective channel protein 1 (VDAC1) purification using, 344–345
 for yeast cell lysis, 458
 Triton X-100 solution for purification of VDAC1 from mitochondria (recipe), 348
 TroponinC family, binding parameters of, 52t
 TRP (transient receptor potential) channels, 277, 278
 Trypan blue, 161
 Trypsin solution for HEK-293 cells (recipe), 275
 Tsiens, Robert, 157
 TTX (tetrodotoxin), 144, 423, 439
 Two-dimensional imaging of fast intracellular Ca^{2+} release (protocol), 130–133
 discussion, 131, 133
 materials, 130
 method, 131, 132f
 recipe, 133
 troubleshooting, 131
 Two-photon fluorescence, 2–3, 2f
 Two-photon microscopy
 calcium signaling investigations using, 113–138
 multibeam two-photon imaging of fast Ca^{2+} signals in Langendorff mouse heart (protocol), 121–125
 discussion, 124
 materials, 121–122
 methods, 122–124, 123f
 recipes, 124
 troubleshooting, 124
 overview, 13–17
 auto-fluorescence, 16
 basic elements of, 14f
 excitation wavelength selection, 15
 implications for design, 14
 laser damage, 14
 photobleaching, 15
 second-harmonic imaging, 16
 trans-illumination imaging, 16–17, 17f
 Two-photon photolysis combined with kilobeam array scanner to probe calcium signaling in cardiomyocytes (protocol), 116–120
 discussion, 119
 materials, 116–117
 method, 117–118, 118f
 recipes, 119–120
 troubleshooting, 118–119
 Two-pore channels, 531
 Tyrode's external solution (recipe), 437, 443
 Tyrode solution for bath perfusion (recipe), 418, 424, 429
 Tyrode solution for probing Ca^{2+} signaling (recipe), 120, 124, 129, 133, 138
- U**
- Ubiquitin promoter, 474
UBQ10 promoter, 474
 Ultrasonicators, 572–573
 Uneven illumination, 34–35, 36f, 49
 Unfolded-protein response (UPR), 242
- V**
- Vacuole, yeast, 445–451, 446f, 449f, 451f
 Vapor-phase chloride, 485
 Vcx1 protein/gene, 446f, 447, 448t, 449, 449f
 VDAC1. *See* Voltage-dependent anion-selective channel protein 1
 VDCC (voltage-dependent Ca^{2+} channel), 555–556
 Vegetable oil (EGTA-washed) (recipe), 204
 Verapamil, 447
 Veratridine, 405
 VGCC (voltage-gated Ca^{2+} channel), 139–140, 142
 Video synchronization, in combined transmitted and fluorescence microscopy, 7
 Vision suite (software), 82, 86
 Vitronectin, 401
 Voltage-dependent anion-selective channel protein 1 (VDAC1), 337–341, 338f
 assay of Ca^{2+} transport by VDAC1 reconstituted into liposomes, 355–358
 discussion, 357
 materials, 355–356
 method, 356–357
 Ca^{2+} efflux from VDAC1-containing proteoliposomes, 356–357
 liposome preparation, 356
 reconstitution of VDAC1 and loading with Ca^{2+} , 356
 recipe, 357
 troubleshooting, 357
 channel conductance and permeability, 340
 conformational states, 340
 permeability to Ca^{2+} , 340–341, 353
 purification from rat liver mitochondria (protocol), 343–348
 discussion, 346
 materials, 343–344
 method, 344–346
 isolation of mitochondria, 344
 VDAC1 purification using carboxymethyl (CM)-cellulose, 345–346
 VDAC1 purification using LDAO, 345
 VDAC1 purification using reactive red-agarose (RRA), 345
 VDAC1 purification using Triton X-100, 344–345
 recipes, 346–348
 troubleshooting, 346
 reconstitution of purified VDAC1 into lipid bilayer and recording of channel conductance (protocol), 349–354

- Voltage-dependent anion-selective channel protein 1 (VDAC1) (*Continued*)
 discussion, 353
 materials, 349–350
 method, 350–353, 351f
 assembly of bilayer system, 350–351
 formation of planar bilayer membrane, 351
 reconstitution of VDAC1 into planar lipid bilayer, 351–352
 recording currents through VDAC1, 352–353, 352f
 recipes, 353
 troubleshooting, 353
- Voltage-dependent Ca^{2+} channel (VDCC), 555–556
- Voltage-gated Ca^{2+} channel (VGCC), 139–140, 142
- Voltage-sensitive Ca^{2+} channels, patch-clamp recordings and, 257–276
 overview, 257–262
 whole-cell recordings of Ca^{2+} currents from isolated neonatal dorsal root ganglion (DRG) neurons (protocol), 264–270
 discussion, 268–269
 materials, 264–265
 method, 265–267
 recipes, 269–270
 troubleshooting, 267–268
- whole-cell recordings of recombinant channels heterologously expressed in HEK-293 cells (protocol), 271–275, 273f
 discussion, 274
 materials, 271–272
 method, 272–274, 273f
 recipes, 275
 troubleshooting, 274
- W**
- Wash buffer for Ca^{2+} -uptake assay (recipe), 254
- Waste, disposal of, 573, 574
- Western blot
 rapid western blotting protocol *Xenopus* oocyte, 395–398
 for SERCA expression, 239, 239f, 250–251, 250f
- Wide-field fluorescence microscopy
 combined transmitted and fluorescence microscopy, 7
 overview, 5–7
 advantages of, 6
 disadvantages of, 6–7
 image quality, 6
 imaging ions, 6
 LED light sources, 5
 sample selection for, 6
- Wsc1p/Slg1p, 450
- X**
- Xenopus*
 calcium transients in developing *Xenopus* spinal neurons (protocol), 399–409
 discussion, 407–408
 materials, 399–400
 method, 400–407
 data analysis, 404
 general imaging methodology, 401–402, 402f
 imaging filopodial transients in vivo, 403–404, 403f
 imaging growth cone transients in vivo, 402–403, 403f
 imposition of Ca^{2+} spikes, 405, 405f
 manipulation of Ca^{2+} dynamics in growth cones, 406
 manipulation of filopodial Ca^{2+} dynamics, 406–407
 perturbation experiments, 404
 preparation of dissociated cell cultures, 400–401
 preparation of spinal cords, 401
 recipes, 408–409
 troubleshooting, 407
- oocyte as single-cell model for studying Ca^{2+} signaling, 381–398
 advantages of, 382–385, 383t, 384t
 nuclear microinjection to assess impact of heterologously expressed proteins (protocol), 388–394
 discussion, 392
 materials, 388–389
 method, 389–391
 recipes, 393
 troubleshooting, 391–392
- overview, 381–382
 rapid western blotting protocol, 395–398
 discussion, 397–398, 397f
 materials, 395–396
 method, 396–397
 recipe, 398
- XPPAUT (software), 562–564, 564f
- Y**
- Yeast
 calcium (Ca^{2+}) signaling, 445–464
 calcium homeostasis, 445–448
 calcium transport, 446–448, 448t
 measuring calcium concentration, 452
 measuring calcium uptake using ^{45}Ca (protocol), 463–464
 materials, 463
 method, 463–464
 monitoring yeast intracellular Ca^{2+} levels using in vivo bioluminescence assay (protocol), 456–459
 materials, 456–457
 method, 457–459, 458f
 troubleshooting, 459
- signaling pathways, 448–452
 calcium-binding proteins, 448–450, 449f
 glucose-induced signaling, 450–451, 451f
 intracellular calcium modification, 450–452, 451f
 programmed cell death, 451–452
 total cellular Ca^{2+} measurements using flame photometry (protocol), 460–462
 materials, 460–461
 method, 461–462
 troubleshooting, 462
- Saccharomyces* Genome Database (SGD), 445
- Yellow Cameleon indicators
 advantages of, 467
 binding parameters of, 52t
 for Ca^{2+} imaging in plants
 high-resolution imaging of cytoplasmic Ca^{2+} dynamics in *Arabidopsis* roots (protocol), 474–478
 data analysis, 477
 discussion, 477
 imaging, 475–477
 materials, 474–475
 method, 475–477, 476f
 plant growth, 475
 sample preparation, 475, 476f
 troubleshooting, 477
- imaging of mitochondrial and nuclear Ca^{2+} dynamics in *Arabidopsis* roots (protocol), 484–488
 CLSM analysis, 486
 discussion, 487
 imaging analyses, 486
 materials, 484–485
 method, 485–486
 recipes, 487
 sample preparation, 486
 seed sterilization and growth of plantlets, 485–486
 troubleshooting, 487
- live cell imaging of cytoplasmic and nuclear Ca^{2+} dynamics in *Arabidopsis* roots (protocol), 479–483
 Ca^{2+} imaging, 480
 data analyses, 482
 discussion, 482
 materials, 479–480
 method, 480–482, 481f
 plant growth, 480
 recipes, 483
 sample preparation, 480, 481f
 troubleshooting, 482
- live cell imaging of cytoplasmic Ca^{2+} dynamics in *Arabidopsis* guard cells (protocol), 469–473
 data analysis, 472
 discussion, 473
 imaging, 471–472
 materials, 469–470
 method, 470–472, 471f
 plant growth, 470
 recipe, 473
 sample preparation, 470–471, 471f
 troubleshooting, 472–473
- overview, 466–467, 466f
 Förster resonance energy transfer (FRET), 466–467, 466f, 472, 477
 functional principle of, 466f
 Yvc1 protein/gene, 446, 446f, 447, 448t, 449f, 450–451
- Z**
- Zebrafish embryos, aequorin introduction into, 193–210, 195f
 microinjecting holo-aequorin into dechorionated and intact zebrafish embryos (protocol), 197–205
 dechorionating embryos, 200
 discussion, 203
 imaging chamber preparation, 198–200, 200f
 materials, 197–198
 method, 198–202
 micropipette preparation, 198, 199f
 recipes, 205
 troubleshooting, 202–203
- reconstitution of holo-aequorin with apoaequorin mRNA and coelenterazine in zebrafish embryos (protocol), 206–210
 discussion, 209
 materials, 206–207
 method, 207–209
 recipes, 210
 troubleshooting, 209